

Recombinant Human PPP3R1 Protein (His Tag)

Catalog Number: PKSH030576

Note: *Centrifuge before opening to ensure complete recovery of vial contents.*


Description

Species	Human
Source	Baculovirus-Insect Cells-derived Human PPP3R1 protein Gly2-Val170, with an N-terminal His
Calculated MW	21.4 kDa
Observed MW	20 kDa
Accession	P63098
Bio-activity	Using the Octet RED System, the affinity constant (Kd) of human PPP3R1-His bound to Human PPIA-His was 6 nM.

Properties

Purity	> 87 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile 50mM Tris, 100mM NaCl, pH 8.0, 10% glycerol, 2mM DTT Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 87 % as determined by reducing SDS-PAGE.

Background

For Research Use Only

Toll-free: 1-888-852-8623
Web: www.elabscience.com

Tel: 1-832-243-6086
Email: techsupport@elabscience.com

Fax: 1-832-243-6017

PPP3R1 belongs to the calcineurin regulatory subunit family. It is a regulatory subunit of calcineurin. Calcineurin is composed of two subunits: calcineurin A (CnA) and calcineurin B (CnB). Dephosphorylation of the nuclear factor of activated T-cells (NF-AT) by Calcineurin is essential for NF-AT activation; nuclear translocation; and early gene expression in T-cells. PPP3R1 is a Ser/Thr-specific calcium and calmodulin-dependent protein phosphatase which takes a vital part in the T cell activation pathway. PPP3R1 is involved in protein dephosphorylation; NFAT protein import into nucleus (ortholog) and epithelial to mesenchymal transition (ortholog). It participates in calcineurin signaling pathway; mitogen activated protein kinase signaling pathway. PPP3R1 interacts with (+)-pilocarpine; 2;4-dinitrotoluene and ammonium chloride. It contains four EF-hand domains and four functional calcium-binding sites. PPP3R1 play an improtant role in the T cell activation pathway.

For Research Use Only

Toll-free: 1-888-852-8623
Web: www.elabscience.com

Tel: 1-832-243-6086
Email: techsupport@elabscience.com

Fax: 1-832-243-6017