Elabscience®

Histone H3 Polyclonal Antibody

catalog number: E-AB-31686

Note: Centrifuge before opening to ensure complete recovery of vial contents.

| Description | |
|----------------------|--|
| Reactivity | Human;Mouse;Rat |
| Immunogen | Synthesized peptide derived from human Histone H3 around the non-phosphorylation |
| | site of Ser10. |
| Host | Rabbit |
| Isotype | IgG |
| Purification | Affinity purification |
| Buffer | Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer, 0.5% protein |
| | protectant and 50% glycerol. |
| Applications | Recommended Dilution |
| WB | 1:500-1:2000 |
| IHC | 1:100-1:300 |
| IF | 1:200-1:1000 |
| Data | |
| | MCF-7 |
| Western Blot analys | sis of MCF7 cells using Histone H3 |
| • | ntibody at dilution of 1:2000. |
| Obs | erved-MW:20 kDa |
| Calc | ulated-MW:15 kDa |
| Preparation & Storag | je |
| Storage | Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles. |
| Shipping | The product is shipped with ice pack, upon receipt, store it immediately at the |
| | temperature recommended. |
| Background | |
| | - 1 N 1 |

Core component of nucleosome. Nucleosomes wrap and compact DNA into chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a template. Histones thereby play a central role in transcription regulation, DNA repair, DNA replication and chromosomal stability. DNA accessibility is regulated via a complex set of post-translational modifications of histones, also called histone code, and nucleosome remodeling.

For Research Use Only