# **Elabscience Biotechnology Co., Ltd.**



A Reliable Research Partner in Life Science and Medicine

# AF/LE Purified Anti-Human CD80 Antibody[2D10]

catalog number: E-AB-F12320

Note: Centrifuge before opening to ensure complete recovery of vial contents.

#### Description

Reactivity Human

Immunogen Recombinant Human CD80 protein

Host Mouse

**Isotype** Mouse IgG1, κ

Clone 2D10

**Purification** >98%, Protein A/G purified

Conjugation None (AF/LE)


Buffer Sterile PBS, pH 7.2. < 1.0 EU per mg of the antibody as determined by the LAL

method.

## Applications Recommended Dilution

FCM  $2 \mu g/mL(0.5 \times 10^6 - 1 \times 10^6 \text{ cells})$ 

#### Data



Raji were stained with 0.2 μg AF/LE Purified Anti-Human CD80 Antibody[2D10] (Right) and 0.2 μg Mouse IgG1, κ

Isotype Control (Left), followed by Elab Fluor® 647-conjugated Goat Anti-Mouse IgG Secondary Antibody.

#### **Preparation & Storage**

Storage Storage, avoid freeze /

thaw cycles. This preparation contains no preservatives, thus it should be handled

under aseptic conditions.

Shipping Ice bag

### **Background**

B7-1 and B7-2, together with their receptors CD28 and CTLA-4, constitute one of the dominant co-stimulatory pathways that regulate T- and B-cell responses. Although both CTLA-4 and CD28 can bind to the same ligands, CTL A-4 binds to B7-1 and B7-2 with a 20-100 fold higher affinity than CD28 and is involved in the down-regulation of the immune response. B7-1 is expressed on activated B cells, activated T cells, and macrophages.