# **SCP2 Polyclonal Antibody**

catalog number: E-AB-52211



Note: Centrifuge before opening to ensure complete recovery of vial contents.

## Description

Reactivity Human; Mouse; Rat

**Immunogen** Fusion protein of human SCP2

Host Rabbit
Isotype IgG

**Purification** Antigen affinity purification


**Conjugation** Unconjugated

**buffer** Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol.

Applications	Recommended Dilution
WB	1:500-1:2000

IHC 1:25-1:100

#### Data



Western blot analysis of Mouse testis tissue lysate using SCP2 Polyclonal Antibody at dilution of 1:350

Immunohistochemistry of paraffin-embedded Human liver cancer tissue using SCP2 Polyclonal Antibody at dilution of  $1:35(\times 200)$ 

Observed-MV: Refer to figures
Calculated-MV: 59 kDa

#### Preparation & Storage

Storage Storage Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.

**Shipping** The product is shipped with ice pack, upon receipt, store it immediately at the

temperature recommended.

#### Background

This gene encodes two proteins: sterol carrier protein X (SCPx) and sterol carrier protein 2 (SCP2), as a result of transcription initiation from 2 independently regulated promoters. The transcript initiated from the proximal promoter encodes the longer SCPx protein, and the transcript initiated from the distal promoter encodes the shorter SCP2 protein, with the 2 proteins sharing a common C-terminus. Evidence suggests that the SCPx protein is a peroxisome-associated thiolase that is involved in the oxidation of branched chain fatty acids, while the SCP2 protein is thought to be an intracellular lipid transfer protein. This gene is highly expressed in organs involved in lipid metabolism, and may play a role in Zellweger syndrome, in which cells are deficient in peroxisomes and have impaired bile acid synthesis. Alternative splicing of this gene produces multiple transcript variants, some encoding different isoforms.

### For Research Use Only