

A Reliable Research Partner in Life Science and Medicine

PE/Cyanine7 Anti-Mouse CD51 Antibody[RMV-7]

Catalog Number: E-AB-F1235UH

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity Mouse
Host Rat

Isotype Rat IgG1, κ
Clone No. RMV-7

Isotype Control PE/Cyanine7 Rat IgG1, κ Isotype Control[HRPN] [Product E-AB-F09823H]

Conjugation PE/Cyanine 7

Conjugation Information PE/Cyanine7 is designed to be excited by the Blue (488 nm), Green (532 nm) and

yellow-green (561 nm) lasers and detected using an optical filter centered near 775 nm

(e.g., a 780/60 nm bandpass filter).

Storage Buffer Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer.

Applications Recommended usage


FCM Each lot of this antibody is quality control tested by flow cytometric analysis. Please

check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use. We suggest each investigator should titrate the

reagent to obtain optimal results [The recommended concentration is 0.1-1 μg/10⁶ cells

in 100 µL volume].

Data



C57BL/6 murine bone marrow cells are stained with PE/Cyanine7 Anti-Mouse CD51 Antibody (filled gray histogram). Unstained bone marrow cells (empty black histogram) are used as control.

Preparation & Storage

Storage Keep as concentrated solution.

This product can be stored at 2-8°C for 12 months. Please protected from prolonged

exposure to light and do not freeze.

Shipping Ice bag

Antigen Information

Alternate Names ITGAV;Integrin alpha-V;Integrin aV chain;Vitronectin Receptor;aV integrin

 Uniprot ID
 P43406

 Gene ID
 16410

For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017

Web:www.elabscience.com Email:techsupport@elabscience.com

Elabscience Bionovation Inc.

A Reliable Research Partner in Life Science and Medicine

Background

Elabscience®

CD51 is a 140 kD protein, also known as αV integrin, vitronectin receptor, and integrin αV . It is a member of the integrin family, expressed on activated T cells, polymorphonuclear granulocytes, platelets, blastocysts, and osteoclasts. CD51 forms heterodimers by association with integrins $\beta 1$, $\beta 3$, $\beta 5$ or $\beta 6$; these complexes then act as receptors for multiple extracellular matrix proteins (ECM). The αV integrin heterodimers have varied functions in development, stimulation/activation and homeostasis. The primary ligands for CD51 complexes are fibronectin, fibrinogen, vitronectin, thrombspondin, von Willebrand factor, and CD31. The RMV-7 antibody has been reported to block binding of CD51 to vitronectin, fibronectin, and CD31 in some cell types, as well as blocking LAK cell cytotoxicity.

 Toll-free: 1-888-852-8623
 Tel: 1-832-243-6086
 Fax: 1-832-243-6017

 Web:www.elabscience.com
 Email:techsupport@elabscience.com