


Recombinant Calmodulin 1/2/3 Monoclonal Antibody

catalog number: AN301731L

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description	
Reactivity	Human;Mouse;Rat
Immunogen	Recombinant human Calmodulin 1/2/3 fragment
Host	Rabbit
Isotype	Rabbit IgG
Clone	C131353
Purification	Protein A purified
Buffer	PBS, 50% glycerol, 0.05% Proclin 300, 0.05% protein protectant.
Applications	
Recommended Dilution	
WB	1:1000-1:5000

Data



Western Blot with Calmodulin 1/2/3 Monoclonal Antibody at dilution of 1:1000. Lane 1: HeLa, Lane 2: NIH/3T3

Observed-MW:17 kDa
Calculated-MW:17 kDa



Western Blot with Calmodulin 1/2/3 Monoclonal Antibody at dilution of 1:5000. Lane 1: Mouse brain, Lane 2: Rat brain

Observed-MW:17 kDa
Calculated-MW:17 kDa

Preparation & Storage

Storage	Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.
Shipping	Ice bag

Background

Calmodulin mediates the control of a large number of enzymes, ion channels, aquaporins and other proteins through calcium-binding. Calmodulin (CaM) is a ubiquitously expressed Ca²⁺ binding protein best known for its ability to activate CaM-dependent kinases in a variety of cells. It was initially identified in its role as an activator of cAMP PDE. In fact, calmodulin was first known as the calcium-dependent regulator protein of PDE. Calmodulin is involved, directly and indirectly, in modulation of several ion channels, including Na⁺, K⁺, L-type Ca²⁺, and RYR2 channels. The main binding partner of calmodulin in cardiac cells is RyR2, which regulates Ca²⁺ release from the sarcoplasmic reticulum. Acting as an intracellular Ca²⁺ sensor, Ca²⁺-saturated calmodulin binds RYR2 to inhibit Ca²⁺ release by stabilizing the RyR2 channel closed state.

For Research Use Only

Toll-free: 1-888-852-8623
Web: www.elabscience.com

Tel: 1-832-243-6086
Email: techsupport@elabscience.com

Fax: 1-832-243-6017

Rev. V1.2