

M-CSF/CSF-1/MGI-IM, Human, Recombinant**Cat. No. : GPCK042****产品信息**

物种	Human
表达宿主	E.coli
序列信息	MEEVSEYCSHMIGSGHLQSLQRLIDSQMETSCQITFEFVDQEQLKDPVCYLKKAFLLVQDIME DTMRFRDNTNPNAIAIVQLQELSLRLKSCFTKDYEEDKACVRTFYETPLQLLEKVKNVFNETKN LLDKDWNIFSKNCNNSFAECSSQGHERQSEGS with polyhistidine tag at the C-terminus.
检索号	P09603.2
标签	His-tag at the C-terminus
分子量	13.34 kDa
有效期	12 months
生物活性	Measure by its ability to induce proliferation in NFS-60 cells. The ED50 for this effect is < 1 ng/mL. The specific activity of recombinant human M-CSF is approximately > 2.5 × 10 ⁸ IU/mg.

产品特性

内毒素 (EU/μg)	< 0.1
保存	Lyophilized protein should be stored at -5~-20°C for 1 year. Upon reconstitution, store at 2-8°C for up to 1 week. Further dilute in a buffer containing a carrier protein or stabilizer (e.g. 0.1% BSA, 10% FBS, 5% HSA or 5% trehalose solution), protein aliquots should be stored at -5~-20°C or -80°C for 3-6 months.
运输	Ambient temperature or ice pack.
制剂	The protein was lyophilized from a 0.2 μm filtered solution containing 1 × PBS, pH 7.4.
复溶	It is recommended to reconstitute the lyophilized protein in sterile water to a concentration not less than 100 μg/mL. Do Not Vortex! Vigorous shaking may impair the biological activity of the protein.

背景介绍

Macrophage Colony-Stimulating Factor (M-CSF), is a secreted cytokine which causes hematopoietic stem cells to differentiate into macrophages or other related cell types. The active form of M-CSF/CSF-1 is found extracellularly as a disulfide-linked homodimer, and is thought to be produced by proteolytic cleavage of membrane-bound precursors. M-CSF/CSF-1 induces cells of the monocyte/macrophage lineage. It also plays a role in immunological defenses, bone metabolism, lipoproteins clearance, fertility and pregnancy. Upregulation of M-CSF/CSF-1 in the infarcted myocardium may have an active role in healing not only through its effects on cells of monocyte/macrophage lineage, but also by regulating endothelial cell chemokine expression.

网站: www.procell.com.cn

电话: 400-999-2100

邮箱: techsupport@procell.com.cn

地址: 湖北省武汉市高新大道858号生物医药产业园三期C4栋

