

## Recombinant Mouse DKK3 protein (His Tag)

Catalog Number: PDMM100207

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


### Description

|                      |   |
|----------------------|---|
| <b>Species</b>       | Mouse   |
| <b>Source</b>        | HEK293 Cells-derived Mouse DKK3 protein Met1-Ile349, with an C-terminal His |
| <b>Calculated MW</b> | 38.3 kDa  |
| <b>Observed MW</b>   | 60 kDa  |
| <b>Accession</b>     | Q9QUN9  |
| <b>Bio-activity</b>  | Not validated for activity  |

### Properties

|                       |  |
|-----------------------|--|
| <b>Purity</b>         | > 95% as determined by reducing SDS-PAGE.  |
| <b>Endotoxin</b>      | < 1.0 EU/mg of the protein as determined by the LAL method   |
| <b>Storage</b>        | Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months. |
| <b>Shipping</b>       | This product is provided as lyophilized powder which is shipped with ice packs.  |
| <b>Formulation</b>    | Lyophilized from a 0.2 µm filtered solution in PBS with 5% Trehalose and 5% Mannitol.  |
| <b>Reconstitution</b> | It is recommended that sterile water be added to the vial to prepare a stock solution of 0.5 mg/mL. Concentration is measured by UV-Vis.   |

### Data



> 95 % as determined by reducing SDS-PAGE.

### Background

DKK3 (dickkopf related protein 3) is a member of the dickkopf-related family consisting of DKK1, DKK2, DKK3 and DKK4. It is a secreted protein, and also known as REIC (Reduced Expansion in Immortalized Cells). The DKK3 protein is proposed to function as a secreted tumor suppressor since it is downregulated in a number of cancer cells and prostate cancer tissue and may be a promising candidate molecule for therapeutic interference. DKK3 protein is also a negative regulator of beta-catenin and its downregulation contribute to an activation of the beta-catenin signaling pathway.

### For Research Use Only