



A Reliable Research Partner in Life Science and Medicine

## PE/Cyanine 5.5 Anti-Human CD14 Antibody [M5E2]

Catalog Number: E-AB-F1209I

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity Human Host Mouse

**Isotype** Mouse IgG2a, κ

Clone No. M5E2

**Isotype Control** PE/Cyanine5.5 Mouse IgG2a, κ Isotype Control[C1.18.4] [Product E-AB-F09802I]

Conjugation PE/Cyanine 5.5

Conjugation Information PE/Cyanine5.5 is designed to be excited by the Blue (488 nm), Green (532 nm) and

yellow-green (561 nm) lasers and detected using an optical filter centered near 690 nm

(e.g., a 690/50 nm bandpass filter).

Storage Buffer Phosphate buffered solution, pH 7.2, containing 0.09% sodium azide and 1% BSA.

Applications Recommended usage

FCM Each lot of this antibody is quality control tested by flow cytometric analysis. The amount

of the reagent is suggested to be used 5  $\mu$ L of antibody per test (million cells in 100  $\mu$ L staining volume or per 100  $\mu$ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for

individual use.

**Preparation & Storage** 

**Storage** Keep as concentrated solution.

This product can be stored at 2-8°C for 12 months. Please protected from prolonged

exposure to light and do not freeze.

Shipping Ice bag

**Antigen Information** 

Alternate Names CD14;Monocyte differentiation antigen CD14;Myeloid cell-specific leucine-rich

glycoprotein

 Uniprot ID
 P08571

 Gene ID
 929

Background CD14 is a 53-55 kD glycosylphosphatidylinositol (GPI)-linked membrane glycoprotein

Web: www.elabscience.cn

also known as LPS receptor. CD14 is expressed at high levels on monocytes and macrophages, and at lower levels on granulocytes. Some dendritic cell populations such as interfollicular dendritic cells, reticular dendritic cells, and Langerhans cells have also been reported to express CD14. As a high-affinity receptor for LPS, CD14 is involved in the clearance of gram-negative pathogens, and in the upregulation of adhesion molecules and expression of cytokines in monocytes and neutrophils.

## For Research Use Only