



A Reliable Research Partner in Life Science and Medicine

AF/LE Purified Anti-Human CD8a Antibody[OKT-8]

catalog number: E-AB-F11100

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity Human

Immunogen Recombinant Human CD8 protein

Host Mouse

Isotype Mouse IgG2a, κ

Clone OKT-8

Purification >98%, Protein A/G purified


Buffer Sterile PBS, pH 7.2. < 1.0 EU per mg of the antibody as determined by the LAL

method.

Applications	Recommended Dilution

FCM $2 \mu g/mL(0.5 \times 10^6 - 1 \times 10^6 \text{ cells})$

Data



Human peripheral blood lymphocytes were stained with 0.2 μg AF/LE Purified Anti-Human CD8a Antibody[OKT-8] (Right) and 0.2 μg Mouse IgG2a, κ Isotype Control (Left),

followed by Elab Fluor® 647-conjugated Goat Anti-Mouse IgG Secondary Antibody.

Preparation & Storage

Storage Store at 4°C valid for 12 months or -20°C valid for long term storage, avoid freeze /

thaw cycles. This preparation contains no preservatives, thus it should be handled

under aseptic conditions.

Shipping lce bag

Background

CD8a is a 32-34 kD type I glycoprotein. It forms a homodimer (CD8a/a) or heterodimer (CD8a/b) with CD8b. CD8, also known as T8 and Leu2, is a member of the immunoglobulin superfamily found on the majority of thymocytes, a subset of peripheral blood T cells, and NK cells (which express almost exclusively CD8a homodimers). CD8 acts as a co-receptor with MHC class I-restricted T cell receptors in antigen recognition and T cell activation and has been shown to play a role in thymic differentiation. Two domains in CD8a are important for function: the extracellular IgSF domain binds the α3 domain of MHC class I and the cytoplasmic CXCP motif binds the tyrosine kinase p56 Lck.

For Research Use Only

 Toll-free: 1-888-852-8623
 Tel: 1-832-243-6086
 Fax: 1-832-243-6017

 Web: www.elabscience.com
 Email: techsupport@elabscience.com
 Rev. V2.6