

## Recombinant Human SCGN/Secretagoin Protein (Human Cells, His Tag)

Catalog Number: PKSH033536

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


### Description

<b>Species</b>	Human
<b>Source</b>	HEK293 Cells-derived Human SCGN/Secretagoin protein Met1-Pro276, with an C-terminal His
<b>Calculated MW</b>	33.1 kDa
<b>Observed MW</b>	32 kDa
<b>Accession</b>	O76038
<b>Bio-activity</b>	Not validated for activity

### Properties

<b>Purity</b>	> 95 % as determined by reducing SDS-PAGE.
<b>Endotoxin</b>	< 1.0 EU per µg of the protein as determined by the LAL method.
<b>Storage</b>	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
<b>Shipping</b>	This product is provided as lyophilized powder which is shipped with ice packs.
<b>Formulation</b>	Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4. Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization.
<b>Reconstitution</b>	Please refer to the specific buffer information in the printed manual.

### Data



> 95 % as determined by reducing SDS-PAGE.

### Background

Secretagoin (SCGN) is a secreted calcium-binding protein that is found in the cytoplasm; a small proportion is associated with secretory granules and membrane fractions. SCGN contains six EF-hand domains, related to calbindin D-28K and calretinin. SCGN is thought to be involved in KCL-stimulated calcium flux and cell proliferation. SCGN can be detected in human serum after ischemic neuronal damage. SCGN may function to negatively control growth and differentiation rates and, thus, indirectly inhibit cell replication.

### For Research Use Only