Elabscience®

HIRIP3 Polyclonal Antibody


catalog number: E-AB-52761


Note: Centrifuge before opening to ensure complete recovery of vial contents.

| Description | |
|--------------|--|
| Reactivity | Human |
| Immunogen | Fusion protein of human HIRIP3 |
| Host | Rabbit |
| Isotype | IgG |
| Purification | Antigen affinity purification |
| Buffer | Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol. |
| | |

| Applications | Recommended Dilution |
|--------------|----------------------|
| IHC | 1:50-1:300 |

Data





Immunohistochemistry of paraffin-embedded Human esophagus cancer tissue using HIRIP3 Polyclonal Antibody at dilution of 1:60(×200)

Immunohistochemistry of paraffin-embedded Human thyroid cancer tissue using HIRIP3 Polyclonal Antibody at dilution of 1:60(×200)

| Preparation & Storage | |
|-----------------------|--|
| Storage | Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles. |
| Shipping | The product is shipped with ice pack, upon receipt, store it immediately at the temperature recommended. |

Background

The HIRA protein shares sequence similarity with Hir1p and Hir2p, the two corepressors of histone gene transcription characterized in the yeast, Saccharomyces cerevisiae. The structural features of the HIRA protein suggest that it may function as part of a multiprotein complex. Several cDNAs encoding HIRA-interacting proteins, or HIRIPs, have been identified. In vitro, the protein encoded by this gene binds HIRA, as well as H2B and H3 core histones, indicating that a complex containing HIRA-HIRIP3 could function in some aspects of chromatin and histone metabolism. Alternatively spliced transcript variants encoding distinct isoforms have been found for this gene.