

Recombinant Human ADGRE5 Protein(His Tag)

Catalog Number: PDMH100241

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Species	Human
Source	Mammalian-derived Human ADGRE5 protein Gln21-Leu530, with an C-terminal His
Calculated MW	55.9 kDa
Observed MW	80-100 kDa
Accession	P48960
Bio-activity	Not validated for activity

Properties

Purity	> 90% as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU/mg of the protein as determined by the LAL method
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from a 0.2 µm filtered solution in PBS with 5% Trehalose and 5% Mannitol.
Reconstitution	It is recommended that sterile water be added to the vial to prepare a stock solution of 0.5 mg/mL. Concentration is measured by UV-Vis.

Data



SDS-PAGE analysis of Human ADGRE5 proteins, 2 µg/lane of Recombinant Human ADGRE5 proteins was resolved with an SDS-PAGE under reducing conditions, showing bands at 55.9KD

Background

This gene encodes a member of the EGF-TM7 subfamily of adhesion G protein-coupled receptors, which mediate cell-cell interactions. These proteins are cleaved by self-catalytic proteolysis into a large extracellular subunit and seven-span transmembrane subunit, which associate at the cell surface as a receptor complex. The encoded protein may play a role in cell adhesion as well as leukocyte recruitment, activation and migration, and contains multiple extracellular EGF-like repeats which mediate binding to chondroitin sulfate and the cell surface complement regulatory protein CD55. Expression of this gene may play a role in the progression of several types of cancer. Alternatively spliced transcript variants encoding multiple isoforms with an 3 to 5 EGF-like repeats have been observed for this gene. This gene is found in a cluster with an other EGF-TM7 genes on the short arm of chromosome 19.