


(KO Validated) NT5E/CD73 Polyclonal Antibody

catalog number: E-AB-60564

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description	
Reactivity	Human;Mouse;Rat
Immunogen	Recombinant fusion protein of human NT5E / CD73
Host	Rabbit
Isotype	Rabbit IgG
Purification	Affinity purification
Buffer	Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol.
Applications	Recommended Dilution
WB	1:500-1:2000
IHC	1:50-1:200
IF	1:50-1:200

Data




Western blot analysis of extracts of various cell lines using CD73/NT5E Polyclonal Antibody at 1:1000 dilution.

Observed-MW:70 kDa
Calculated-MW:57 kDa/63 kDa





Western blot analysis of extracts of Rat eye using CD73/NT5E Polyclonal Antibody at 1:1000 dilution.

Observed-MW:70 kDa
Calculated-MW:57 kDa/63 kDa




Western blot analysis of extracts from normal (control) and CD73/NT5E Polyclonal antibody knockout (KO) HCT116 cells using CD73/NT5E Polyclonal antibody at 1:3000 dilution.

Observed-MW:70 kDa
Calculated-MW:57 kDa/63 kDa



Immunohistochemistry of paraffin-embedded human liver cancer using [KO Validated] CD73/NT5E Polyclonal antibody at dilution of 1:50 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.



For Research Use Only


Toll-free: 1-888-852-8623
Web: www.elabscience.com

Tel: 1-832-243-6086
Email: techsupport@elabscience.com

Fax: 1-832-243-6017

Rev. V1.8

Immunohistochemistry of paraffin-embedded human tonsil using [KO Validated] CD73/NT5E Polyclonal antibody at dilution of 1:50 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.



Immunohistochemistry of paraffin-embedded human tonsil using [KO Validated] CD73/NT5E Polyclonal antibody at dilution of 1:50 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.

Immunohistochemistry of paraffin-embedded human liver cancer using [KO Validated] CD73/NT5E Polyclonal antibody at dilution of 1:50 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.

Preparation & Storage

Storage

Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.

Shipping

The product is shipped with ice pack, upon receipt, store it immediately at the temperature recommended.

Background

The protein encoded by this gene is a plasma membrane protein that catalyzes the conversion of extracellular nucleotides to membrane-permeable nucleosides. The encoded protein is used as a determinant of lymphocyte differentiation. Defects in this gene can lead to the calcification of joints and arteries. Two transcript variants encoding different isoforms have been found for this gene.

For Research Use Only

Toll-free: 1-888-852-8623
Web: www.elabscience.com

Tel: 1-832-243-6086
Email: techsupport@elabscience.com

Fax: 1-832-243-6017

Rev. V1.8