

Osteoprotegerin/TNFRSF11B Monoclonal Antibody(Detector)

catalog number: **AN001770P**

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Reactivity	Human
Immunogen	Recombinant Human Osteoprotegerin/TNFRSF11B protein expressed by Mammalian
Host	Rat
Isotype	Rat IgG1
Clone	7D9
Purification	Protein A/G Purification
Buffer	Phosphate buffered solution, pH 7.2, containing 0.05% Proclin300.

Applications Recommended Dilution

ELISA Detector	0.1-0.4 µg/mL
-----------------------	---------------

Data



Sandwich ELISA-Recombinant Human Osteoprotegerin/TNFRSF11B protein standard curve. Background subtracted standard curve using Osteoprotegerin/TNFRSF11B antibody(AN001760P) (Capture), Osteoprotegerin/TNFRSF11B Antibody(AN001770P)(Detector) in sandwich ELISA. The reference range value for Recombinant Human Osteoprotegerin/TNFRSF11B protein is 0.16-10 ng/mL.

Preparation & Storage

Storage	Store at 4°C valid for 12 months or -20°C valid for long term storage, avoid freeze / thaw cycles.
Shipping	The product is shipped with ice pack, upon receipt, store it immediately at the temperature recommended.

Background

For Research Use Only

TNFRSF11B is a secreted protein; containing 2 death domains and 4 TNFR-Cys repeats. TNFRSF11B is a decoy receptor for the receptor activator of nuclear factor kappa B ligand (RANKL). By binding RANKL, TNFRSF11B inhibits nuclear kappa B (NF-κB) which is a central and rapid acting transcription factor for immune-related genes; and a key regulator of inflammation; innate immunity; and cell survival and differentiation. TNFRSF11B levels are influenced by voltage-dependent calcium channels Cav1.2. TNFRSF11B can reduce the production of osteoclasts by inhibiting the differentiation of osteoclast precursors (osteoclasts are related to monocytes/macrophages and are derived from granulocyte/macrophage-forming colony units (CFU-GM)) into osteoclasts and also regulates the resorption of osteoclasts in vitro and in vivo. TNFRSF11B binding to RANKL on osteoblast/stromal cells; blocks the RANKL-RANK ligand interaction between