

Recombinant Human Profilin-2/PFN2 Protein

Catalog Number: PKSH032935

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

| | |
|----------------------|---|
| Species | Human |
| Source | E.coli-derived Human Profilin-2;PFN2 protein Met 1-Phe140 |
| Calculated MW | 15.0 kDa |
| Observed MW | 14 kDa |
| Accession | P35080 |
| Bio-activity | Not validated for activity |

Properties

| | |
|-----------------------|---|
| Purity | > 95 % as determined by reducing SDS-PAGE. |
| Endotoxin | < 1.0 EU per µg of the protein as determined by the LAL method. |
| Storage | Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months. |
| Shipping | This product is provided as lyophilized powder which is shipped with ice packs. |
| Formulation | Lyophilized from a 0.2 µm filtered solution of 20mM Tris-HCl, 150mM NaCl, pH 8.0. Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. |
| Reconstitution | Please refer to the specific buffer information in the printed manual. Please refer to the printed manual for detailed information. |

Data



> 95 % as determined by reducing SDS-PAGE.

Background

Profilin-II (PFN2) is ubiquitous protein which belongs to the profilin family. PFN2 binds to actin, then affects the structure of the cytoskeleton. At high concentrations, profiling prevents the polymerization of actin, while increases that at low concentrations. PFN2 is a ubiquitous actin monomer-binding protein. It regulates actin polymerization in response to extra cellular signals. PFN2 binds to PIP2; it inhibits the formation of IP3 and DG.

For Research Use Only

Toll-free: 1-888-852-8623
Web: www.elabscience.com

Tel: 1-832-243-6086
Email: techsupport@elabscience.com

Fax: 1-832-243-6017