Purified Anti-Human CD127 Antibody[A019D5]

catalog number: E-AB-F11520P




Note: Centrifuge before opening to ensure complete recovery of vial contents.

| Description | |
|--------------|--|
| Reactivity | Human |
| Immunogen | Recombinant Human CD127 protein |
| Host | Mouse |
| Is otype | Mouse IgGl, ĸ |
| Clone | A019D5 |
| Purification | >98%, Protein A/G purified |
| Conjugation | Unconjugated |
| buffer | PBS, pH 7.2. Contains 0.05% proclin 300. |
| Applications | Recommended Dilution |

 $2 \,\mu g/mL(1 \times 10^5 - 5 \times 10^5 \text{ cells})$

FCM

Data



Human peripheral blood lymphocytes were stained with 0.2 μg Purified Anti-Human CD127 Antibody[A019D5] (Right) and 0.2 μg Mouse IgG1, κ Isotype Control (Left), followed by Alexa Fluor® 647-conjugated Goat Anti-Mouse IgG

Secondary Antibody.

| Preparation & Storage | |
|-----------------------|---|
| Storage | Store at 4°C valid for 12 months or -20°C valid for long term storage, avoid freeze / |
| | thaw cycles. |
| Shipping | Ice bag |

Background

The protein encoded by this gene is a receptor for interleukin 7 (IL7). The function of this receptor requires the interleukin 2 receptor, gamma chain (IL2RG), which is a common gamma chain shared by the receptors of various cytokines, including interleukins 2, 4, 7, 9, and 15. This protein has been shown to play a critical role in V(D)J recombination during lymphocyte development. Defects in this gene may be associated with severe combined immunodeficiency (SCID). Alternatively spliced transcript variants have been found.

For Research Use Only