

CCL16/HCC-4 Polyclonal Antibody(Capture/Detector)

catalog number: AN004640P

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

| | |
|---------------------|---|
| Reactivity | Human |
| Immunogen | Recombinant Human CCL16/HCC-4 Protein expressed by E.coli |
| Host | Rabbit |
| Isotype | Rabbit IgG |
| Purification | Antigen Affinity Purification |
| Buffer | Phosphate buffered solution, pH 7.2, containing 0.05% Proclin300. |

Applications Recommended Dilution

| | |
|-----------------------|---------------|
| ELISA Capture | 2-8 µg/mL |
| ELISA Detector | 0.1-0.4 µg/mL |

Data



Sandwich ELISA-Recombinant Human CCL16/HCC-4
Protein standard curve. Background subtracted standard curve
using Anti-CCL16/HCC-4 antibody(AN004640P)
(Capture), Anti-CCL16/HCC-4 antibody(AN004640P)
(Detector). The reference range value is 0.16~10 ng/mL for
human.

Preparation & Storage

| | |
|-----------------|--|
| Storage | Store at 4°C valid for 12 months or -20°C valid for long term storage, avoid freeze / thaw cycles. |
| Shipping | The product is shipped with ice pack, upon receipt, store it immediately at the temperature recommended. |

Background

This gene is one of several cytokine genes clustered on the q-arm of chromosome 17. Cytokines are a family of secreted proteins involved in immunoregulatory and inflammatory processes. The CC cytokines are proteins characterized by two adjacent cysteines. The cytokine encoded by this gene displays chemotactic activity for lymphocytes and monocytes but not for neutrophils. This cytokine also shows a potent myelosuppressive activity and suppresses proliferation of myeloid progenitor cells. The expression of this gene is upregulated by IL-10.

For Research Use Only