## **Elabscience**®


## Elab Fluor<sup>®</sup> 488 Anti-Human CD73 Antibody[AD2]

Catalog Number: E-AB-F1242L

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description	
Reactivity	Human
Host	Mouse
lsotype	Mouse IgG1, ĸ
Clone No.	AD2
Isotype Control	Elab Fluor <sup>®</sup> 488 Mouse IgG1, κ Isotype Control[MOPC-21] [Product E-AB-F09792L]
Conjugation	Elab Fluor <sup>®</sup> 488
Conjugation Information	Elab Fluor <sup>®</sup> 488 is designed to be excited by the Blue laser (488 nm) and detected using an optical filter centered near 520 nm (e.g., a 525/40 nm bandpass filter).
Storage Buffer	Phosphate buffered solution, pH 7.2, containing 0.09% sodium azide and 1% BSA.
Applications	Recommended usage
FCM	Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5 $\mu$ L of antibody per test (million cells in 100 $\mu$ L staining volume or per 100 $\mu$ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.

Data



Human peripheral blood lymphocytes are stained with APC

Anti-Human CD3 Antibody and Elab Fluor<sup>®</sup> 488 Anti-Human CD73 Antibody[AD2] (Left). Lymphocytes are stained with

APC Anti-Human CD3 Antibody and Elab Fluor  $^{I\!\!R}$  488 Mouse IgG1,  $\kappa$  Isotype Control (Right).

Preparation & Storage	
Storage	Keep as concentrated solution.
	This product can be stored at 2-8°C for 12 months. Please protected from prolonged exposure to light and do not freeze.
Shipping	Ice bag
Antigen Information	
Alternate Names	5'-NTE.C3.1.3.5L-VAP-2NT5E;Ecto-5'-nucleotidase
Uniprot ID	P21589
Gene ID	4907

## For Research Use Only

## **Elabscience**®

Background

Elabscience Biotechnology Co., Ltd. A Reliable Research Partner in Life Science and Medicine

CD73 is a 70 kD glycophosphatidylinositol (GPI)-linked 5'-nucleotidase, which is also known as ecto-5'-nucleotidase. It converts adenosine monophosphate (AMP) to adenosine. CD73 is expressed on subsets of T and B cells, mesenchymal stem cells, follicular dendritic cells, endothelial cells, and epithelial cells. It has been reported that CD73 costimulates T cell activation, and mediates adhesion of lymphocytes to follicular dendritic cells and endothelial cells.