

A Reliable Research Partner in Life Science and Medicine

# Recombinant CCR1/CD191 Monoclonal Antibody

catalog number: AN300171P

Note: Centrifuge before opening to ensure complete recovery of vial contents.

#### **Description**

Reactivity Human

Immunogen A synthetic peptide corresponding to the center region of the Human CCR1/CD191


HostRabbitIsotypeIgGCloneA1285PurificationProtein A

Buffer 0.2 µm filtered solution in PBS

## Applications Recommended Dilution

WB 1:500-1:2000IP 1-4 μL/mg of lysate

#### Data



ssection Blot with CCR1/CD19

Immunoprecipitation analysis using 2 µL anti-CCR1/CD191 Monoclonal Antibody and 15 µl of 50 % Protein G agarose. Western blot was performed from the immunoprecipitate using CCR1/CD191 Monoclonal Antibody at a dilution of 1:100. Lane A:0.5 mg Hela Whole Cell Lysate, Lane B:0.5 mg 293T Whole Cell Lysate

Observed-MW:43 kDa Calculated-MW:43 kDa

Western Blot with CCR1/CD191 Monoclonal Antibody at dilution of 1:500. Lane A: HEK293 Whole Cell Lysate, Lane B: 293T Whole Cell Lysate, Lysates/proteins at 30 µg per lane.

Observed-MW:43 kDa Calculated-MW:43 kDa

Rev. V1.2

#### **Preparation & Storage**

Storage This antibody can be stored at 2°C-8°C for one month without detectable loss of

activity. Antibody products are stable for twelve months from date of receipt when stored at -20°C to -80°C. Preservative-Free. Avoid repeated freeze-thaw cycles.

Shipping Ice bag

...

**Background** 

# For Research Use Only

 Toll-free: 1-888-852-8623
 Tel: 1-832-243-6086
 Fax: 1-832-243-6017

 Web: www.elabscience.com
 Email: techsupport@elabscience.com

# Elabscience®

### **Elabscience Bionovation Inc.**

A Reliable Research Partner in Life Science and Medicine

This gene encodes a member of the beta chemokine receptor family, which is predicted to be a seven transmembrane protein similar to G protein-coupled receptors. The ligands of this receptor include macrophage inflammatory protein 1 alpha (MIP-1 alpha), regulated on activation normal T expressed and secreted protein (RANTES), monocyte chemoattractant protein 3 (MCP-3), and myeloid progenitor inhibitory factor-1 (MPIF-1). Chemokines and their receptors mediated signal transduction are critical for the recruitment of effector immune cells to the site of inflammation. Knockout studies of the mouse homolog suggested the roles of this gene in host protection from inflammatory response, and susceptibility to virus and parasite. This gene and other chemokine receptor genes, including CCR2, CCRL2, CCR3, CCR5 and CCXCR1, are found to form a gene cluster on chromosome 3p. [provided by RefSeq, Jul 2005]

For Research Use Only

 Toll-free: 1-888-852-8623
 Tel: 1-832-243-6086
 Fax: 1-832-243-6017

 Web: www.elabscience.com
 Email: techsupport@elabscience.com
 Rev. V1.2