Elabscience®


GPM6A Polyclonal Antibody

catalog number: E-AB-63925


Note: Centrifuge before opening to ensure complete recovery of vial contents.

| Description | |
|--------------|--|
| Reactivity | Human;Mouse |
| Immunogen | A synthetic peptide of human GPM6A (NP_005268.1). |
| Host | Rabbit |
| Isotype | IgG |
| Purification | Affinity purification |
| Buffer | Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol. |
| Applications | Recommended Dilution |
| WB | 1:500-1:2000 |
| IHC | 1:50-1:200 |

Data



Observed-MW:31 kDa



Immunohistochemistry of paraffin-embedded Human stomach using GPM6A Polyclonal Antibody at dilution of

1:100 (40x lens).

| Preparation & Storage | |
|-----------------------|--|
| Storage | Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles. |
| Shipping | The product is shipped with ice pack,upon receipt, store it immediately at the |
| | temperature recommended. |

Background

For Research Use Only

Immunohistochemistry of paraffin-embedded Mouse stomach using GPM6A Polyclonal Antibody at dilution of 1:100 (40x lens).

Western blot analysis of extracts of U-87MG cells using GPM6A Polyclonal Antibody at dilution of 1:1000.

Calculated-MW:29 kDa/30 kDa/31 kDa

Elabscience®

Neuronal membrane glycoprotein M6-ais aproteinthat in humans is encoded by theGPM6Agene. Involved in neuronal differentiation, including differentiation and migration of neuronal stem cells. Plays a role in neuronal plasticity and is involved in neurite and filopodia outgrowth, filopodia motility and probably synapse formation. GPM6A-induced filopodia formation involves mitogen-activated protein kinase (MAPK) and Src signaling pathways. May be involved in neuronal NGF-dependent Ca2+ influx. May be involved in regulation of endocytosis and intracellular trafficking of G-protein-coupled receptors (GPCRs); enhances internalization and recycling of mu-type opioid receptor.

For Research Use Only