

**FGF-17 (C-6His), Human, Recombinant**

Cat. No. : PCK130

**General Information**

<b>Synonyms</b>	Fibroblast Growth Factor 17;FGF-17;FGF17
<b>Species</b>	Human
<b>Expression host</b>	Human Cells
<b>Sequence</b>	Thr23-Thr216
<b>Accession</b>	O60258
<b>Tag</b>	C-6His
<b>Mol mass</b>	22.6 kDa
<b>Expiration date</b>	12 months
<b>Bio activity</b>	Measured in a cell proliferation assay using Balb/3T3 mouse embryonic fibroblast cells. The ED50 for this effect is 2.1 µg/mL.

**Product feature**

<b>Purity</b>	> 95% as determined by reducing SDS-PAGE.
<b>Endotoxin (EU/µg)</b>	< 0.1
<b>Storage</b>	Lyophilized protein should be stored at -5~-20°C, stable for one year after receipt. Reconstituted protein solution can be stored at 2-8°C for 2-7 days. Aliquots of reconstituted samples are stable at -5~-20°C for 3 months.
<b>Shipping</b>	Ice bag
<b>Formulation</b>	Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4.
<b>Reconstitution</b>	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 µg/mL. Dissolve the lyophilized protein in sterile water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

**Background**

Fibroblast Growth Factor 17 (FGF17) is a member of the heparin-binding Growth Factors family that is prominently expressed in the cerebellum and cortex. Proteins of this family possess broad mitogenic and cell survival activities and they are involved in a variety of biological processes including embryonic development cell growth, morphogenesis, tissue repair, tumor growth, and invasion. FGF17 plays an important role in the regulation of embryonic development and it acts as signaling molecule in the induction and patterning of the embryonic brain. In addition, FGF17 stimulates the proliferation and activation of cells that express FGF Receptors.