

A Reliable Research Partner in Life Science and Medicine

PE Anti-Mouse CD48 Antibody [HM48-1]

Catalog Number: E-AB-F1017UD

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity Mouse

Host Armenian Hamster Isotype Armenian Hamster IgG

Clone No. HM48-1

PE Armenian Hamster IgG Isotype Control[PIP] [Product E-AB-F09853D] Isotype Control

Conjugation

Conjugation Information PE is designed to be excited by the Blue (488 nm), Green (532 nm) and Yellow-Green

(561 nm) lasers and detected using an optical filter centered near 575 nm (e.g., a 585/42

nm bandpass filter).

Storage Buffer Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer.

Applications Recommended usage


FCM Each lot of this antibody is quality control tested by flow cytometric analysis. Please

> check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use. We suggest each investigator should titrate the

> reagent to obtain optimal results [The recommended concentration is 0.1-1 μg/10⁶ cells

in 100 µL volume].

Data



C57BL/6 murine splenocytes are stained with PE Anti-Mouse CD48 Antibody (filled gray histogram). Unstained splenocytes (empty black histogram) are used as control.

Preparation & Storage

Storage Keep as concentrated solution.

This product can be stored at 2-8°C for 12 months. Please protected from prolonged

exposure to light and do not freeze.

Shipping Ice bag

Antigen Information

Alternate Names BCM1 surface antigen;BLAST-1;CD48;CD48 antigen;Cd48;HM48-1;MRC OX-45

surface antigen; SLAMF2; sgp-60

Uniprot ID P18181 Gene ID 12506

For Research Use Only

Fax: 1-832-243-6017 Tel: 1-832-243-6086 Toll-free: 1-888-852-8623 Email:techsupport@elabscience.com

Web:www.elabscience.com

Elabscience Bionovation Inc.



A Reliable Research Partner in Life Science and Medicine

Background

CD48 is a 45 kD GPl-anchored glycoprotein also known as BCM1, Blast-1 (human), and OX-45 (rat). It is a member of the lg superfamily, expressed on T and B cells and monocytes/macrophages. It plays a role in adhesion and T cell recognition. The primary ligands for CD48 are CD2 and CD244.

Fax: 1-832-243-6017