

# Recombinant Human LAIR1 Protein (Fc Tag)

Catalog Number:PKSH030465



**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


## Description

<b>Synonyms</b>	Leukocyte-Associated Immunoglobulin-Like Receptor 1;LAIR-1;hLAIR1;CD305;LAIR1
<b>Species</b>	Human
<b>Expression Host</b>	HEK293 Cells
<b>Sequence</b>	Met 1-Tyr165
<b>Accession</b>	Q6GTX8.1
<b>Calculated Molecular Weight</b>	42.8 kDa
<b>Tag</b>	C-hFc

## Properties

<b>Purity</b>	> 95 % as determined by reducing SDS-PAGE.
<b>Endotoxin</b>	< 1.0 EU per µg of the protein as determined by the LAL method.
<b>Storage</b>	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
<b>Shipping</b>	This product is provided as lyophilized powder which is shipped with ice packs.
<b>Formulation</b>	Lyophilized from sterile PBS, pH 7.4 Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
<b>Reconstitution</b>	Please refer to the printed manual for detailed information.

## Data



> 95 % as determined by reducing SDS-PAGE.

## Background

Leukocyte associated Ig-like receptor-1 (LAIR1) is a surface molecule expressed on human mononuclear leukocytes that functions as an inhibitory receptor on human NK cells. In addition to NK cells, LAIR1 is expressed on T cells, B cells, macrophages, and dendritic cells. It is predicted to mediate inhibitory functions based on the presence of immunoreceptor tyrosine-based inhibitory motifs (ITIMs) in its cytoplasmic domain. Cross-linking of LAIR1 on human T cell clones results in inhibition of cytotoxicity only in T cell clones that lack CD28 and are able to spontaneously lyse certain targets in vitro. Moreover, the cytolytic activity of freshly isolated T cells, which is thought to be mainly due to "effector" T cells, can be inhibited by anti-LAIR1 mAb. Thus, LAIR1 functions as an inhibitory receptor not only on NK cells, but also on human T cells. This indicates that LAIR1 provides a mechanism of regulation of effector T cells and may play a role in the inhibition of unwanted bystander responses mediated by Ag-specific T cells.

## For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Web: [www.elabscience.com](http://www.elabscience.com)

Tel: 1-832-243-6086

Email: [techsupport@elabscience.com](mailto:techsupport@elabscience.com)

Fax: 1-832-243-6017

# Recombinant Human LAIR1 Protein (Fc Tag)

Catalog Number:PKSH030465



## For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Web: [www.elabscience.com](http://www.elabscience.com)

Tel: 1-832-243-6086

Email: [techsupport@elabscience.com](mailto:techsupport@elabscience.com)

Fax: 1-832-243-6017