

Recombinant Mouse TGFBR3/Betaglycan Protein (His Tag)

Catalog Number: PKSM040623

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Species	Mouse
Source	HEK293 Cells-derived Mouse TGFBR3/Betaglycan protein Met 1-Arg 399, with an C-terminal His
Calculated MW	86.3 kDa
Observed MW	80-90 kDa
Accession	NP_072075.2
Bio-activity	Not validated for activity

Properties

Purity	> 95 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile PBS, pH 7.4 Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

For Research Use Only

Betaglycan also known as transforming growth factor beta receptor III (TGFBR3), is a cell-surface chondroitin sulfate / heparan sulfate proteoglycan. TGFBR3 is a transforming growth factor (TGF)-beta type III receptor. This receptor is a membrane proteoglycan that often functions as a co-receptor with other TGF-beta receptor superfamily members. Ectodomain shedding produces soluble TGFBR3, which may inhibit TGFβ signaling. Decreased expression of this receptor has been observed in various cancers. TGFBR3 is the TGF-β component most commonly downregulated among localized human prostate cancer studies. TGFBR3 knockdown led to focus formation and enhanced expression of CD133, a marker found on prostate cancer stem cells. TGFBR3 is an accessory receptor that binds to and modulates the activities of both transforming growth factor-beta (TGFβ;) and inhibin, two members of the TGFβ; superfamily of growth factors that regulate many aspects of reproductive biology. TGFBR3 is known to be expressed in adult testis and ovary, but little is known about this receptor during gonadogenesis.

For Research Use Only

Toll-free: 1-888-852-8623
Web: www.elabscience.com

Tel: 1-832-243-6086
Email: techsupport@elabscience.com

Fax: 1-832-243-6017