

A Reliable Research Partner in Life Science and Medicine

Elab Fluor® 488 Anti-Mouse/Human CD11b Antibody[M1/70]

Catalog Number: E-AB-F1081L

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity Human; Mouse

Host Rat

IsotypeRat lgG2b, κ Clone No.M1/70

Isotype Control Elab Fluor® 488 Rat IgG2b, κ Isotype Control[LTF-2] [Product E-AB-F09842L]

Conjugation Elab Fluor® 488

Conjugation Information Elab Fluor® 488 is designed to be excited by the Blue laser (488 nm) and detected using

an optical filter centered near 520 nm (e.g., a 525/40 nm bandpass filter).


Storage Buffer Phosphate buffered solution, pH 7.2, containing 0.09% sodium azide and 1% BSA.

Applications Recommended usage

FCM

Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5 μ L of antibody per test (million cells in 100 μ L staining volume or per 100 μ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.

Data



C57BL/6 murine bone marrow cells are stained with Elab

Fluor[®] 488 Anti-Mouse/Human CD11b Antibody (filled gray histogram). Unstained bone marrow cells (empty black histogram) are used as control.

Preparation & Storage

Storage Keep as concentrated solution.

This product can be stored at 2-8°C for 12 months. Please protected from prolonged

exposure to light and do not freeze.

Shipping Ice bag

Antigen Information

Alternate Names CD11 antigen-like family member B;CD11b;CR-3 alpha chain;Integrin alpha-M;Itgam;

Web: www.elabscience.cn

Leukocyte adhesion receptor MO1

 Uniprot ID
 P05555;P11215

 Gene ID
 16409,3684

For Research Use Only



Elabscience Biotechnology Co., Ltd.

A Reliable Research Partner in Life Science and Medicine

Background

CD11b is a 170 kD glycoprotein also known as αM integrin, Mac-1 α subunit, Mol, CR3, and Ly-40. CD11b is a member of the integrin family, primarily expressed on granulocytes, monocytes/macrophages, dendritic cells, NK cells, and subsets of T and B cells. CD11b non-covalently associates with CD18 (β2 integrin) to form Mac-1. Mac-1 plays an important role in cell-cell interaction by binding its ligands ICAM-1 (CD54), ICAM-2 (CD102), ICAM-4 (CD242), iC3b, and fibrinogen.

Web: www.elabscience.cn