

A Reliable Research Partner in Life Science and Medicine

Claudin18.2 Monoclonal Antibody

catalog number: AN200149P

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity Human

Immunogen Human Claudin18.2 mRNA

 Host
 Mouse

 Isotype
 IgG2a

 Clone
 A1001

 Purification
 Protein A

Buffer 0.2 µm filtered solution in PBS


Applications Recommended Dilution

WB 1:500-1:2000 IHC-P 1:500-1:2000


Data




Immunohistochemistry of paraffin-embedded human stomach using Claudin18.2 Monoclonal Antibody at dilution of 1:1000.



Immunohistochemistry of paraffin-embedded human gastric cancer using Claudin18.2 Monoclonal Antibody at dilution of 1:1000.



Immunohistochemistry of paraffin-embedded human lung using Claudin18.2 Monoclonal Antibody at dilution of 1:1000.



Western Blot with Claudin18.2 Monoclonal Antibody at dilution of 1:1000. Lane A: Human Claudin18.2-VLP (Full Length) Protein 500ng, Lane B: Virus-Like Particle (VLP) isotype control Protein 500ng

Rev. V1.2

Preparation & Storage

Storage This antibody can be stored at 2°C-8°C for one month without detectable loss of

activity. Antibody products are stable for twelve months from date of receipt when stored at -20°C to -80°C. Preservative-Free. Avoid repeated freeze-thaw cycles.

Shipping Ice bag

Background

For Research Use Only

 Toll-free: 1-888-852-8623
 Tel: 1-832-243-6086
 Fax: 1-832-243-6017

 Web: www.elabscience.com
 Email: techsupport@elabscience.com

Elabscience®

Elabscience Bionovation Inc.

A Reliable Research Partner in Life Science and Medicine

This gene encodes a member of the claudin family. Claudins are integral membrane proteins and components of tight junction strands. Tight junction strands serve as a physical barrier to prevent solutes and water from passing freely through the paracellular space between epithelial or endothelial cell sheets, and also play critical roles in maintaining cell polarity and signal transductions. This gene is upregulated in patients with ulcerative colitis and highly overexpressed in infiltrating ductal adenocarcinomas. PKC/MAPK/AP-1 (protein kinase C/mitogen-activated protein kinase/activator protein-2) dependent pathway regulates the expression of this gene in gastric cells. Alternatively spliced transcript variants encoding different isoforms have been identified.

For Research Use Only

 Toll-free: 1-888-852-8623
 Tel: 1-832-243-6086
 Fax: 1-832-243-6017

 Web: www.elabscience.com
 Email: techsupport@elabscience.com
 Rev. V1.2