



A Reliable Research Partner in Life Science and Medicine

# **Human TIMP-1 Antibody Pair Set**

Catalog No. E-KAB-0067 Applications ELISA

**Synonyms** TIMP1, CLGI, EPA, EPO, HCI

## **Kit components & Storage**

Title	Specifications	Storage
Human TIMP-1 Capture Antibody	1 vial, 100 μ g	Store at -20°C for one year.
		Avoid freeze / thaw cycles.
Human TIMP-1 Detection Antibody	1 vial, 50 μL	Store at -20°C for one year.
(Biotin)		Avoid freeze / thaw cycles.

Note: Centrifuge before opening to ensure complete recovery of vial contents.

#### **Product Information**

Items		Characteristic (E-KAB-0067)	
		Human TIMP-1 Capture Antibody	Human TIMP-1 Detection Antibody
			(Biotin)
Immunogen	Immunogen	Recombinant Human TIMP-1 protein	Recombinant Human TIMP-1 protein
Information	Swissprot	P01033	
Product details	Reactivity	Human	Human
	Host	Mouse	Goat
	Conjugation	Unconjugated	Biotin
	Concentration	0.5mg/mL	/
	Buffer	PBS with 0.04% Proclin 300, 50%	PBS with 0.04% Proclin 300, 1%
		glycerol, pH 7.4	protective protein, 50% glycerol, pH
			7.4
	Purify	Protein A or G	Antigen Affinity
	Specificity	Detects Human TIMP-1 in ELISAs.	

For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017 Web: <a href="mailto:www.elabscience.com">www.elabscience.com</a> Email: <a href="mailto:techsupport@elabscience.com">techsupport@elabscience.com</a>





A Reliable Research Partner in Life Science and Medicine

### **Applications**

Human TIMP-1 Sandwich ELISA Assay:

	Recommended	Reagent	Images
	Concentration/Dilution		
ELISA	0.5-4µg/mL	Human TIMP-1 Capture Antibody	
Capture			Aigs 1
ELISA	1:1000-1:10000	Human TIMP-1 Detection	Optical Density
Detection		Antibody (Biotin)	0.01 0.01 1 10 100  Human TIMP-1 concentration(ng/mL)

Note: This standard curve is only for demonstration purposes. A standard curve should be generated for each assay!

#### **Background**

This gene belongs to the TIMP gene family. The proteins encoded by this gene family are natural inhibitors of the matrix metalloproteinases (MMPs), a group of peptidases involved in degradation of the extracellular matrix. In addition to its inhibitory role against most of the known MMPs, the encoded protein is able to promote cell proliferation in a wide range of cell types, and may also have an anti-apoptotic function. Transcription of this gene is highly inducible in response to many cytokines and hormones. In addition, the expression from some but not all inactive X chromosomes suggests that this gene inactivation is polymorphic in human females. This gene is located within intron 6 of the synapsin I gene and is transcribed in the opposite direction.