

# Recombinant Human NEIL1 Protein (His Tag)

Catalog Number:PKSH030794



**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


## Description

<b>Synonyms</b>	FPG1;hFPG1;NEI1
<b>Species</b>	Human
<b>Expression Host</b>	E.coli
<b>Sequence</b>	Met 1-Ser 390
<b>Accession</b>	AAH10876.1
<b>Calculated Molecular Weight</b>	45 kDa
<b>Observed molecular weight</b>	45 kDa
<b>Tag</b>	C-His

## Properties

<b>Purity</b>	> 84 % as determined by reducing SDS-PAGE.
<b>Endotoxin</b>	Please contact us for more information.
<b>Storage</b>	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
<b>Shipping</b>	This product is provided as lyophilized powder which is shipped with ice packs.
<b>Formulation</b>	Lyophilized from sterile 50mM Tris, 150mM NaCl, pH 8.0 Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
<b>Reconstitution</b>	Please refer to the printed manual for detailed information.

## Data



> 84 % as determined by reducing SDS-PAGE.

## Background

NEIL1 is a member of DNA glycosylases. DNA glycosylases are a family homologous to the bacterial Fpg/Nei family. They play a role in base excision repair which is the mechanism by which damaged bases in DNA are removed and replaced. The first step of this process is catalyzed by DNA glycosylases. They remove the damaged nitrogenous base while leaving the sugar-phosphate backbone intact, creating an apurinic/apyrimidinic site, commonly referred to as an AP site. NEIL1 functions in base excision repair of DNA damaged by oxidation or by mutagenic agents. It acts as DNA glycosylase that recognizes and removes damaged bases. NEIL1 prefers to oxidized pyrimidines, such as thymine glycol, formamidopyrimidine (Fapy) and 5-hydroxyuracil. Has marginal activity towards 8-oxoguanine. It has AP (apurinic/apyrimidinic) lyase activity and introduces nicks in the DNA strand and cleaves the DNA backbone by beta-

## For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Web: [www.elabscience.com](http://www.elabscience.com)

Tel: 1-832-243-6086

Email: [techsupport@elabscience.com](mailto:techsupport@elabscience.com)

Fax: 1-832-243-6017

# Recombinant Human NEIL1 Protein (His Tag)

Catalog Number:PKSH030794



delta elimination to generate a single-strand break at the site of the removed base with both 3'- and 5'-phosphates.

---

## For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Web: [www.elabscience.com](http://www.elabscience.com)

Tel: 1-832-243-6086

Email: [techsupport@elabscience.com](mailto:techsupport@elabscience.com)

Fax: 1-832-243-6017