

## Recombinant Human CEACAM7/CGM2 Protein (His Tag)

**Catalog Number:** PKSH032238

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


### Description

<b>Species</b>	Human
<b>Source</b>	HEK293 Cells-derived Human CEACAM7;CGM2 protein Thr36-Phe142, with an C-terminal His
<b>Calculated MW</b>	13.1 kDa
<b>Observed MW</b>	20-30 kDa
<b>Accession</b>	AAI21133.1
<b>Bio-activity</b>	Not validated for activity

### Properties

<b>Purity</b>	> 95 % as determined by reducing SDS-PAGE.
<b>Endotoxin</b>	< 1.0 EU per µg of the protein as determined by the LAL method.
<b>Storage</b>	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
<b>Shipping</b>	This product is provided as lyophilized powder which is shipped with ice packs.
<b>Formulation</b>	Lyophilized from a 0.2 µm filtered solution of 20mM PB, 150mM NaCl, pH7.4. Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization.
	Please refer to the specific buffer information in the printed manual.
<b>Reconstitution</b>	Please refer to the printed manual for detailed information.

### Data



> 95 % as determined by reducing SDS-PAGE.

### Background

Carcinoembryonic Antigen-Related Cell Adhesion Molecule 7 (CEACAM7) is a member of the immunoglobulin superfamily A and CEA family. CEACAM7 localizes to the cell membrane and contains one Ig-like C2-type domain and one Ig-like V-type domain. The expression of CEACAM7 is significantly decreased in rectal cancer. Differences in CEACAM7 expression levels between long-term survivors and those with recurrent disease introduce a potential tumor marker to define a subset of patients who benefit most from adjuvant therapy.

### For Research Use Only