

DRAK2 Polyclonal Antibody

catalog number: E-AB-14744

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Reactivity	Human;Mouse;Rat
Immunogen	Recombinant protein of human STK17B
Host	Rabbit
Isotype	IgG
Purification	Affinity purification
Buffer	Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol.


Applications Recommended Dilution

IHC	1:50-1:200
------------	------------

Data



Immunohistochemistry of paraffin-embedded Human thyroid cancer tissue using DRAK2 Polyclonal Antibody at dilution 1:40



Immunohistochemistry of paraffin-embedded Human liver cancer tissue using DRAK2 Polyclonal Antibody at dilution 1:40

Preparation & Storage

Storage	Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.
Shipping	The product is shipped with ice pack, upon receipt, store it immediately at the temperature recommended.

Background

The full-length STK17B cDNA clone encodes a deduced 372-amino acid protein with a molecular mass of 42.34 kD. The putative kinase domain is located at the N terminus and contains all 11 subdomains conserved among ser/thr kinases. STK17A and STK17B share 59.7% amino acid identity. Northern blot analysis revealed that STK17B was expressed in various tissues, such as heart, placenta, liver, and pancreas, as different-sized transcripts, presumably due to differences in the 3-prime untranslated region. Transient transfection of COS-7 cells showed that STK17B localized in nuclei. Phosphorylates myosin light chains. Acts as a positive regulator of apoptosis.

For Research Use Only