Elab Fluor[®] 647-conjugated Myc-Tag Monoclonal Antibody

catalog number: AN00300M



Note: Centrifuge before opening to ensure complete recovery of vial contents.

-						70		
D	Α		C	11	m	ŤΤ	n	m
\mathbf{L}	·	0	V.	ш	עו	UΙ	v	ш

Reactivity All

Immunogen MYC tag fusion protein

Host Mouse Isotype IgGl Clone 6B10

Purification Protein A/G Purification


Conjugation Elab Fluor® 647(Ex/Em:650nm/665nm)

buffer PBS with 0.05% proclin 300, 1% protective protein and 50% glycerol.pH7.4

Applications Recommended Dilution

FCM $0.2 \mu g \text{ per } 10^6 \text{ cells in a } 100 \mu L \text{ suspension}$

Data



1X10⁶ HEK-293F cells were intracellularly stained with 0.2

ug Elab Fluor[®] 647-conjugated Myc-Tag Monoclonal Antibody(red), or 0.2 ug Elab Fluor[®] 647-conjugated Myc-Tag Monoclonal Antibody Isotype Control (green).

Preparation & Storage

Storage Storage Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles. Protected from

prolonged exposure to light.

Shipping The product is shipped with ice pack, upon receipt, store it immediately at the

temperature recommended.

Background

The MYC tag, derived from the c-MYC protein, is a popular epitope tag for detecting the expression of recombinant proteins in yeast, bacteria, insect, and mammalian cell systems. It can be used for affinity chromatography, then used to separate recombin

For Research Use Only