

Mouse LDLR Antibody Pair Set

Catalog No.	E-KAB-0309	Applications	ELISA
Synonyms	FH, FHC, LDLCQ2, Familial Hypercholesterolemia		

Kit components & Storage

Title	Specifications	Storage
Mouse LDLR Capture Antibody	1 vial, 100 µg	Store at -20°C for one year. Avoid freeze / thaw cycles.
Mouse LDLR Detection Antibody (Biotin)	1 vial, 50 µL	Store at -20°C for one year. Avoid freeze / thaw cycles.


Note: Centrifuge before opening to ensure complete recovery of vial contents.

Product Information

Items		Characteristic (E-KAB-0309)	
		Mouse LDLR Capture Antibody	Mouse LDLR Detection Antibody (Biotin)
Immunogen Information	Immunogen	Recombinant Mouse LDLR protein	Recombinant Mouse LDLR protein
	Swissprot	P35951	
Product details	Reactivity	Mouse	Mouse
	Host	Rabbit	Rabbit
	Conjugation	Unconjugated	Biotin
	Concentration	0.5mg/mL	/
	Buffer	PBS with 0.04% Proclin 300, 50% glycerol, pH 7.4	PBS with 0.04% Proclin 300, 1% protective protein, 50% glycerol, pH 7.4
	Purify	Protein A & Antigen Affinity	Protein A & Antigen Affinity
	Specificity	Detects Mouse LDLR in ELISAs.	

Applications

Mouse LDLR Sandwich ELISA Assay:

	Recommended Concentration/Dilution	Reagent	Images
ELISA Capture	0.5-4μg/mL	Mouse LDLR Capture Antibody	 <p>The graph is a log-log plot. The x-axis is labeled 'Mouse LDLR concentration(ng/mL)' and ranges from 0.01 to 100. The y-axis is labeled 'Optical Density' and ranges from 0.01 to 10. The data points form a smooth, upward-sloping curve, indicating a positive correlation between the concentration of Mouse LDLR and the optical density measured in the assay.</p>
ELISA Detection	1:1000-1:10000	Mouse LDLR Detection Antibody (Biotin)	

Note: This standard curve is only for demonstration purposes. A standard curve should be generated for each assay!

Background

LDLR (low density lipoprotein receptor) is a member of the LDL receptor gene family and is involved in receptor-mediated endocytosis of specific ligands. The LDLR is a cell surface glycoprotein that scavenges LDL from the blood and regulates plasma LDL cholesterol. The cytoplasmic domain of the LDL receptor is necessary for the receptor to cluster in coated pits, which promotes the rapid endocytosis of bound LDL. The protein is highly glycosylated through N- and O-linkages and thus migrates at 100 to 160 kDa bands on SDS-PAGE.