Elabscience®


Elab Fluor[®] Violet 610 Anti-Mouse/Human CD11b Antibody[M1/70]

Catalog Number: E-AB-F1081T

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description	
Reactivity	Human;Mouse
Host	Rat
Isotype	Rat lgG2b, κ
Clone No.	M1/70
Isotype Control	Elab Fluor [®] Violet 610 Rat IgG2b, κ Isotype Control[LTF-2] [Product E-AB-F09842T]
Conjugation	Elab Fluor [®] Violet 610
Conjugation Information	Elab Fluor [®] Violet 610 is designed to be excited by the violet laser (405 nm) and detected using an optical filter centered near 613 nm (e.g., a 615/20 nm bandpass filter).
Storage Buffer	Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant.
Applications	Recommended usage
FCM	Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5 µL of antibody per test (million cells in 100 µL staining volume or per 100 µL of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.

Data



Staining of C57BL/6 murine bone marrow cells with Elab

Fluor[®] Violet 610 Anti-Mouse/Human CD11b Antibody[M1/70] (filled gray histogram) or Elab Fluor[®] Violet 610 Rat IgG2b, κ Isotype Control (empty black histogram).

Total viable cells were used for analysis.

Preparation & Storage	
Storage	Keep as concentrated solution.
	This product can be stored at 2-8°C for 12 months. Please protected from prolonged
	exposure to light and do not freeze.
Shipping	Ice bag
Antigen Information	
Alternate Names	CD11 antigen-like family member B;CD11b;CR-3 alpha chain;Integrin alpha-M;Itgam; Leukocyte adhesion receptor MO1

For Research Use Only

Toll-free: 1-888-852-8623 Web:<u>www.elabscience.com</u>

Elabscience®

Uniprot ID Gene ID Background

P05555;P11215;

16409;3684

CD11b is a 170 kD glycoprotein also known as α M integrin, Mac-1 α subunit, Mol, CR3, and Ly-40. CD11b is a member of the integrin family, primarily expressed on granulocytes, monocytes/macrophages, dendritic cells, NK cells, and subsets of T and B cells. CD11b non-covalently associates with CD18 (β 2 integrin) to form Mac-1. Mac-1 plays an important role in cell-cell interaction by binding its ligands ICAM-1 (CD54), ICAM-2 (CD102), ICAM-4 (CD242), iC3b, and fibrinogen.

For Research Use Only