# **Elabscience**®

## SUPT16H Polyclonal Antibody


### catalog number: E-AB-92122

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description	
Reactivity	Human;Mouse;Rat
Immunogen	Recombinant fusion protein of human SUPT16H
Host	Rabbit
Isotype	IgG
Purification	Affinity purification
Buffer	Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol.
Applications	Recommended Dilution


Аррисацоня	Ke commende a Dilation
WB	1:500-1:2000
IF	1:50-1:200

#### Data




Western blot analysis of extracts of various cell lines using SUPT16H Polyclonal Antibody at1:1000 dilution.

Observed-MW:140 kDa Calculated-MW:119 kDa



Western blot analysis of extracts of Mouse brain using SUPT16H Polyclonal Antibody at 1:1000 dilution.

Observed-MW:140 kDa Calculated-MW:119 kDa



Immunofluorescence analysis of U2OS using SUPT16H Immunofluore Polyclonal Antibody at dilution of 1:100 (40x lens). Blue: Polyclonal An DAPI for nuclear staining.

Immunofluorescence analysis of NIH/3T3 using SUPT16H Polyclonal Antibody at dilution of 1:100 (40x lens). Blue: DAPI for nuclear staining.

Preparation & Storage	
Storage	Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.
Shipping	The product is shipped with ice pack, upon receipt, store it immediately at the
	temperature recommended.

### Background

For Research Use Only

Toll-free: 1-888-852-8623 Web:<u>w w .elabscience.com</u>

Tel: 1-832-243-6086 Email:<u>techsupport@elabscience.com</u>

# **Elabscience**®

Transcription of protein-coding genes can be reconstituted on naked DNA with only the general transcription factors and RNA polymerase II. However, this minimal system cannot transcribe DNA packaged into chromatin, indicating that accessory factors may facilitate access to DNA. One such factor, FACT (facilitates chromatin transcription), interacts specifically with histones H2A/H2B to effect nucleosome disassembly and transcription elongation. FACT is composed of an 80 kDa subunit and a 140 kDa subunit; this gene encodes the 140 kDa subunit.

For Research Use Only