




CCL16 Polyclonal Antibody

catalog number: E-AB-16304

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description	
Reactivity	Human
Immunogen	Synthetic peptide of human CCL16
Host	Rabbit
Isotype	IgG
Purification	Affinity purification
Conjugation	Unconjugated
Buffer	Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol.
Applications	Recommended Dilution
WB	1:500-1:2000
IHC	1:50-1:200

Data	
 <p>Western Blot analysis of HepG2 cell using CCL16 Polyclonal Antibody at dilution of 1:760</p> <p>Calculated-MV:14 kDa</p>	 <p>Immunohistochemistry of paraffin-embedded Human Lymphoma using CCL16 Polyclonal Antibody at dilution of 1:50</p>
 <p>Immunohistochemistry of paraffin-embedded Human brain using CCL16 Polyclonal Antibody at dilution of 1:50</p>	

Preparation & Storage	
Storage	Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.
Shipping	The product is shipped with ice pack, upon receipt, store it immediately at the temperature recommended.

Background	
<p>This gene is one of several cytokine genes clustered on the q-arm of chromosome 17. Cytokines are a family of secreted proteins involved in immunoregulatory and inflammatory processes. The CC cytokines are proteins characterized by two adjacent cysteines. The cytokine encoded by this gene displays chemotactic activity for lymphocytes and monocytes but not for neutrophils. This cytokine also shows a potent myelosuppressive activity and suppresses proliferation of myeloid progenitor cells. The expression of this gene is upregulated by IL-10.</p>	

For Research Use Only