# **Elabscience Biotechnology Co., Ltd.**



A Reliable Research Partner in Life Science and Medicine

# **IDH2 Polyclonal Antibody**

catalog number: E-AB-11319

Note: Centrifuge before opening to ensure complete recovery of vial contents.

#### **Description**

Reactivity Human; Mouse; Rat

Immunogen Recombinant protein of human IDH2

Host Rabbit Isotype IgG


Purification Affinity purification
Conjugation Unconjugated

**Buffer** Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol.


## Applications Recommended Dilution

**WB** 1:500-1:2000 **IHC** 1:100-1:300

#### Data




Western Blot analysis of Human fetal muscle tissue, Jurkat and 293T cell, Hela cell and Mouse liver tissue using IDH2 Polyclonal Antibody at dilution of 1:600



Immunohistochemistry of paraffin-embedded Human cervical cancer using IDH2 Polyclonal Antibody at dilution of 1:60

#### Calculated-MV:51 kDa



Immunohistochemistry of paraffin-embedded Human colon cancer using IDH2 Polyclonal Antibody at dilution of 1:60

#### Preparation & Storage

Storage Storage Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.

**Shipping** The product is shipped with ice pack, upon receipt, store it immediately at the

temperature recommended.

#### Background

#### For Research Use Only

# Elabscience®

### Elabscience Biotechnology Co., Ltd.

A Reliable Research Partner in Life Science and Medicine

Isocitrate dehydrogenases catalyze the oxidative decarboxylation of isocitrate to 2-oxoglutarate. These enzymes belong to two distinct subclasses, one of which utilizes NAD(+) as the electron acceptor and the other NADP(+). Five isocitrate dehydrogenases have been reported: three NAD(+)-dependent isocitrate dehydrogenases, which localize to the mitochondrial matrix, and two NADP(+)-dependent isocitrate dehydrogenases, one of which is mitochondrial and the other predominantly cytosolic. Each NADP(+)-dependent isozyme is a homodimer. The protein encoded by this gene is the NADP(+)-dependent isocitrate dehydrogenase found in the mitochondria. It plays a role in intermediary metabolism and energy production. This protein may tightly associate or interact with the pyruvate dehydrogenase complex. Alternative splicing results in multiple transcript variants.

Web: www.elabscience.cn

For Research Use Only

Tel: 400-999-2100