

Human TRAIL Antibody Pair Set

Catalog No.	E-KAB-0274	Applications	ELISA
Synonyms	APO2L, Apo2-L, TL2, CD253		

Kit components & Storage

Title	Specifications	Storage
Human TRAIL Capture Antibody	1 vial, 100 µg	Store at -20°C for one year. Avoid freeze / thaw cycles.
Human TRAIL Detection Antibody (Biotin)	1 vial, 50 µL	Store at -20°C for one year. Avoid freeze / thaw cycles.

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Product Information

Items		Characteristic (E-KAB-0274)	
		Human TRAIL Capture Antibody	Human TRAIL Detection Antibody (Biotin)
Immunogen Information	Immunogen	Recombinant Human TRAIL protein	Recombinant Human TRAIL protein
	Swissprot	P50591	
Product details	Reactivity	Human	Human
	Host	Goat	Goat
	Conjugation	Unconjugated	Biotin
	Concentration	0.5mg/mL	/
	Buffer	PBS with 0.04% Proclin 300, 50% glycerol, pH 7.4	PBS with 0.04% Proclin 300, 1% protective protein, 50% glycerol, pH 7.4
	Purify	Antigen Affinity	Antigen Affinity
	Specificity	Detects Human TRAIL in ELISAs.	

For Research Use Only

Applications

Human TRAIL Sandwich ELISA Assay:

	Recommended Concentration/Dilution	Reagent	Images
ELISA Capture	0.5-4µg/mL	Human TRAIL Capture Antibody	
ELISA Detection	1:1000-1:10000	Human TRAIL Detection Antibody (Biotin)	

Note: This standard curve is only for demonstration purposes. A standard curve should be generated for each assay!

Background

The protein encoded by this gene is a cytokine that belongs to the tumor necrosis factor (TNF) ligand family. This protein preferentially induces apoptosis in transformed and tumor cells, but does not appear to kill normal cells although it is expressed at a significant level in most normal tissues. This protein binds to several members of TNF receptor superfamily including TNFRSF10A/TRAILR1, TNFRSF10B/TRAILR2, TNFRSF10C/TRAILR3, TNFRSF10D/TRAILR4, and possibly also to TNFRSF11B/OPG. The activity of this protein may be modulated by binding to the decoy receptors TNFRSF10C/TRAILR3, TNFRSF10D/TRAILR4, and TNFRSF11B/OPG that cannot induce apoptosis. The binding of this protein to its receptors has been shown to trigger the activation of MAPK8/JNK, caspase 8, and caspase 3. Alternatively spliced transcript variants encoding different isoforms have been found for this gene.

For Research Use Only