

STRAP Polyclonal Antibody

catalog number: E-AB-92347

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Reactivity	Human;Mouse
Immunogen	Recombinant fusion protein of human STRAP
Host	Rabbit
Isotype	IgG
Purification	Affinity purification
Buffer	Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol.

Applications

Applications	Recommended Dilution
WB	1:500-1:2000
IF	1:50-1:200
IP	1:50-1:200


Data




Western blot analysis of extracts of various cell lines using STRAP Polyclonal Antibody at 1:1000 dilution.

Observed-MW: Refer to figures


Calculated-MW: 38 kDa/39 kDa




Immunofluorescence analysis of H9C2 cells using [KO Validated] STRAP Polyclonal Antibody at dilution of 100 (40x lens). Blue: DAPI for nuclear staining.



Immunofluorescence analysis of L929 cells using [KO Validated] STRAP Polyclonal Antibody at dilution of 100 (40x lens). Blue: DAPI for nuclear staining.



Immunofluorescence analysis of U2OS cells using [KO Validated] STRAP Polyclonal Antibody at dilution of 100 (40x lens). Blue: DAPI for nuclear staining.



For Research Use Only

Immunoprecipitation analysis of 200ug extracts of HeLa cells
using 3ug STRAP Polyclonal Antibody. Western blot was
performed from the immunoprecipitate using STRAP

Polyclonal Antibody at a dilution of 1:1000.

Observed-MW: Refer to figures

Calculated-MW: 38 kDa/39 kDa

Preparation & Storage

Storage Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.

Shipping The product is shipped with ice pack, upon receipt, store it immediately at the temperature recommended.

Background

The SMN complex catalyzes the assembly of small nuclear ribonucleoproteins (snRNPs, the building blocks of the spliceosome, and thereby plays an important role in the splicing of cellular pre-mRNAs. Most spliceosomal snRNPs contain a common set of Sm proteins SNRPB, SNRPD1, SNRPD2, SNRPD3, SNRPE, SNRPF and SNRPG that assemble in a heptameric protein ring on the Sm site of the small nuclear RNA to form the core snRNP (Sm core). In the cytosol, the Sm proteins SNRPD1, SNRPD2, SNRPE, SNRPF and SNRPG are trapped in an inactive 6S pICln-Sm complex by the chaperone CLNS1A that controls the assembly of the core snRNP. To assemble core snRNPs, the SMN complex accepts the trapped 5Sm proteins from CLNS1A forming an intermediate. Binding of snRNA inside 5Sm triggers eviction of the SMN complex, thereby allowing binding of SNRPD3 and SNRPB to complete assembly of the core snRNP. STRAP plays a role in the cellular distribution of the SMN complex. Negatively regulates TGF-beta signaling but positively regulates the PDPK1 kinase activity by enhancing its autophosphorylation and by significantly reducing the association of PDPK1 with 14-3-3 protein.

For Research Use Only