

## Recombinant Human FGF-19/FGF9 Protein (His Tag)

Catalog Number: PKSH032436

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


### Description

<b>Species</b>	Human
<b>Source</b>	E.coli-derived Human FGF-19;FGF9 protein Phe27-Lys216, with an C-terminal His
<b>Calculated MW</b>	22.6 kDa
<b>Observed MW</b>	24 kDa
<b>Accession</b>	O95750
<b>Bio-activity</b>	Measure by its ability to induce 3T3 cells proliferation. The ED <sub>50</sub> for this effect is <51 ng/mL.

### Properties

<b>Purity</b>	> 95 % as determined by reducing SDS-PAGE.
<b>Endotoxin</b>	< 0.1 EU per µg of the protein as determined by the LAL method.
<b>Storage</b>	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
<b>Shipping</b>	This product is provided as lyophilized powder which is shipped with ice packs.
<b>Formulation</b>	Lyophilized from sterile PBS, pH 8.0. Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
<b>Reconstitution</b>	Please refer to the printed manual for detailed information.

### Data



> 95 % as determined by reducing SDS-PAGE.

### Background

Fibroblast growth factor 19 (FGF19) is a secreted protein which belongs to the FGFs family. FGF19 is expressed in fetal brain, cartilage, retina, and adult gall bladder. FGFs modulate cellular activity via at least 5 distinct subfamilies of high-affinity FGF receptors (FGFRs): FGFR-1, -2, -3, and -4, all with intrinsic tyrosine kinase activity. FGFRs can be important for regulation of glucose and lipid homeostasis. FGF19 has important roles as a hormone produced in the ileum in response to bile acid absorption. It has been shown to cause resistance to diet-induced obesity and insulin desensitization and to improve insulin, glucose, and lipid profiles in diabetic rodents. FGF19 can be considered as a regulator of energy expenditure.

### For Research Use Only