



A Reliable Research Partner in Life Science and Medicine

Human OPG Antibody Pair Set

Catalog No. E-KAB-0060 Applications ELISA

Synonyms TNFRSF11B, OCIF, TR1

Kit components & Storage

Title	Specifications	Storage
Human OPG Capture Antibody	1 vial, 100 μ g	Store at -20°C for one year.
		Avoid freeze / thaw cycles.
Human OPG Detection Antibody (Biotin)	1 vial, 50 μL	Store at -20°C for one year.
		Avoid freeze / thaw cycles.

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Product Information

Items		Characteristic (E-KAB-0060)		
		Human OPG Capture Antibody	Human OPG Detection Antibody	
			(Biotin)	
Immunogen	Immunogen	Recombinant Human OPG protein	Recombinant Human OPG protein	
Information	Swissprot	O00300		
Product details	Reactivity	Human	Human	
	Host	Mouse	Goat	
	Conjugation	Unconjugated	Biotin	
	Concentration	0.5mg/mL	/	
	Buffer	PBS with 0.04% Proclin 300, 50%	PBS with 0.04% Proclin 300, 1%	
		glycerol, pH 7.4	protective protein, 50% glycerol, pH	
			7.4	
	Purify	Protein A or G	Antigen Affinity	
	Specificity	Detects Human OPG in ELISAs.	·	

For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017 Web: www.elabscience.com Email: techsupport@elabscience.com





A Reliable Research Partner in Life Science and Medicine

Applications

Human OPG Sandwich ELISA Assay:

	Recommended	Reagent	Images
	Concentration/Dilution		
ELISA	0.5-4µg/mL	Human OPG Capture Antibody	
Capture			Aigu Niga Niga Niga Niga Niga Niga Niga Niga
ELISA Detection	1:1000-1:10000	Human OPG Detection Antibody (Biotin)	Optical Density
Detection		(Biotili)	0.01 0.1 1 10 100 Human OPG concentration(ng/mL)

Note: This standard curve is only for demonstration purposes. A standard curve should be generated for each assay!

Background

The protein encoded by this gene is a member of the TNF-receptor superfamily. This protein is an osteoblast-secreted decoy receptor that functions as a negative regulator of bone resorption. This protein specifically binds to its ligand, osteoprotegerin ligand, both of which are key extracellular regulators of osteoclast development. Studies of the mouse counterpart also suggest that this protein and its ligand play a role in lymph-node organogenesis and vascular calcification. Alternatively spliced transcript variants of this gene have been reported, but their full length nature has not been determined.

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017 Web: www.elabscience.com Email: techsupport@elabscience.com