## **Elabscience**®

## Purified Anti-Human CD105 Antibody[SN6h]

catalog number: AN003640P

Note: Centrifuge before opening to ensure complete recovery of vial contents.

| Description  |   |  |
|--------------|---|--|
| Reactivity   | Human   |  |
| Immunogen    | Recombinant Human CD105 protein   |  |
| Host         | Mouse   |  |
| Isotype      | Mouse IgG1, ĸ   |  |
| Clone        | SN6h  |  |
| Purification | >98%, Protein A/G purified  |  |
| Conjugation  | Unconjugated  |  |
| Buffer       | Phosphate-buffered solution, pH 7.2, containing 0.05% non-protein stabilizer. Dialyze |  |
|              | to completely remove the stabilizer prior to labeling.                                |  |
| Applications | Recommended Dilution  |  |
| FCM          | $2 \ \mu g/mL(1 \times 10^5 - 5 \times 10^5 \text{ cells})$                           |  |
| Data         |   |  |
|              | 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0   |  |

THP-1 were stained with 0.2 μg Purified Anti-Human CD105 Antibody[SN6h] (Right) and 0.2 μg Mouse IgG1, κ Isotype Control (Left), followed by Alexa Fluor® 647-conjugated

Goat Anti-Mouse IgG Secondary Antibody.

| Preparation & Storage |   |
|-----------------------|---|
| Storage               | Store at 4°C valid for 12 months or -20°C valid for long term storage, avoid freeze / |
|                       | thaw cycles.  |
| Shipping              | Ice bag   |
|                       |   |

## Background

Endoglin (CD105) is a 90 kDa type I transmembrane glycoprotein of the zona pellucida (ZP) family of proteins (1-3). Endoglin and Betaglycan/T beta RIII are type III receptors for TGF beta superfamily ligands, sharing 71% aa identity in the transmembrane (TM) and cytoplasmic domains. Endoglin is highly expressed on proliferating vascular endothelial cells, chondrocytes, and syncytiotrophoblasts of term placenta, with lower amounts on hematopoietic, mesenchymal and neural crest stem cells, activated monocytes, and lymphoid and myeloid leukemic cells.