

## PE/Cyanine7 Anti-Human CD123 Antibody[6H6]

Catalog Number: E-AB-F1117H

Note: Centrifuge before opening to ensure complete recovery of vial contents.

### Description


Reactivity	Human
Host	Mouse
Isotype	Mouse IgG1, $\kappa$
Clone No.	6H6
Isotype Control	PE/Cyanine7 Mouse IgG1, $\kappa$ Isotype Control[MOPC-21] [Product E-AB-F09792H]
Conjugation	PE/Cyanine 7
Conjugation Information	PE/Cyanine7 is designed to be excited by the Blue (488 nm), Green (532 nm) and yellow-green (561 nm) lasers and detected using an optical filter centered near 775 nm (e.g., a 780/60 nm bandpass filter).
Storage Buffer	Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer.

### Applications

### Recommended usage

FCM	Each lot of this antibody is quality control tested by flow cytometric analysis. <b>The amount of the reagent is suggested to be used 5 <math>\mu</math>L of antibody per test (million cells in 100 <math>\mu</math>L staining volume or per 100 <math>\mu</math>L of whole blood).</b> Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.
-----	--

### Data



Human peripheral blood mononuclear cells are stained with

Elab Fluor® 488 Anti-Human CD45RO Antibody and PE/Cyanine7 Anti-Human CD123 Antibody[6H6] (Left).

Mononuclear cells are stained with Elab Fluor® 488 Anti-Human CD45RO Antibody and PE/Cyanine7 Mouse IgG1,  $\kappa$  Isotype Control (Right).

### Preparation & Storage

Storage	Keep as concentrated solution. This product can be stored at 2-8°C for 24 months. Please protected from prolonged exposure to light and do not freeze.
Shipping	Ice bag

### Antigen Information

Alternate Names	IL-3 Receptor alpha;IL-3R $\alpha$
-----------------	------------------------------------

### For Research Use Only

**Uniprot ID**

P26951

**Gene ID**

3563

**Background**

CD123 is the 70 kD transmembrane  $\alpha$  chain of the IL-3 receptor. Alone, CD123 binds IL-3 with low affinity; when CD123 associates with CDw131 (common  $\beta$  chain), it binds IL-3 with high affinity. CD123 does not transduce intracellular signals upon binding IL-3 and requires the  $\beta$  chain for this function. CD123 is expressed by myeloid precursors, macrophages, dendritic cells, mast cells, basophils, megakaryocytes, and some B cells.