

Recombinant Podoplanin/PDPN Monoclonal Antibody

catalog number: **AN300498P**

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Reactivity	Mouse
Immunogen	Recombinant Mouse Podoplanin/PDPN Protein
Host	Rabbit
Isotype	IgG
Clone	8C2
Purification	Protein A
Buffer	0.2 µm filtered solution in PBS


Applications Recommended Dilution

IHC-P	1:250-1:1000
--------------	--------------

Data



Immunohistochemistry of paraffin-embedded mouse kidney using Podoplanin/PDPN Monoclonal Antibody at dilution of 1:500.



Immunohistochemistry of paraffin-embedded mouse brain using Podoplanin/PDPN Monoclonal Antibody at dilution of 1:500.

Preparation & Storage

Storage	This antibody can be stored at 2°C-8°C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20°C to -80°C. Preservative-Free. Avoid repeated freeze-thaw cycles.
Shipping	Ice bag

Background

Podoplanin, also known as PDPN, is a type-I integral membrane glycoprotein with diverse distribution in human tissues. The physiological function of this protein may be related to its mucin-type character. The homologous protein in other species has been described as a differentiation antigen and influenza-virus receptor. The specific function of this protein has not been determined. Alternatively spliced transcript variants encoding different isoforms have been identified. PDPN is a mucin-type glycoprotein negatively charged by extensive O-glycosylation and a high content of sialic acid, which expresses the adhesive property. It is selectively expressed in lymphatic endothelium as well as lymphangiomas, Kaposi sarcomas, and in a subset of angiosarcomas with probable lymphatic differentiation. PDPN may contribute to form odontoblastic fiber or function as the anchorage to the tooth development and in proliferating epithelial cells of cervical loop and apical bud. The intensity of podoplanin expression is negatively correlated with the expression of CD34 and factor VIII. Podoplanin would be useful as a diagnostic marker for epithelioid hemangioendothelioma in liver tumors.

For Research Use Only