

## Recombinant Na<sup>+</sup>/K<sup>+</sup> ATPase alpha-1 Monoclonal Antibody

catalog number: **AN301712L**

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.

### Description


<b>Reactivity</b>	Human;Rat;Mouse
<b>Immunogen</b>	Recombinant human Na <sup>+</sup> /K <sup>+</sup> ATPase alpha-1 fragment
<b>Host</b>	Rabbit
<b>Isotype</b>	IgG, κ
<b>Clone</b>	A420
<b>Purification</b>	Protein A purified
<b>Buffer</b>	PBS, 50% glycerol, 0.05% Proclin 300, 0.05% protein protectant.

### Applications

### Recommended Dilution


<b>WB</b>	1:500-1:2000
<b>IHC</b>	1:200-1:1000
<b>IF</b>	1:50

### Data





Western Blot with Na<sup>+</sup>/K<sup>+</sup> ATPase alpha-1 Monoclonal Antibody at dilution of 1:2000. Lane 1: HeLa, Lane 2: RAW264.7, Lane 3: C6, Lane 4: Mouse brain, Lane 5: Rat brain


**Observed-MW:113 kDa**  
**Calculated-MW:113 kDa**



Immunohistochemistry of paraffin-embedded Human kidney using Na<sup>+</sup>/K<sup>+</sup> ATPase alpha-1 Monoclonal Antibody at dilution of 1:1000.



Immunohistochemistry of paraffin-embedded Human cervical cancer using Na<sup>+</sup>/K<sup>+</sup> ATPase alpha-1 Monoclonal Antibody at dilution of 1:1000.



Immunofluorescent analysis of (100% ice-cold methanol) fixed Jurkat cells using anti-Na<sup>+</sup>/K<sup>+</sup> ATPase alpha-1 Monoclonal Antibody at dilution of 1:50.

### For Research Use Only

Toll-free: 1-888-852-8623  
Web: [www.elabscience.com](http://www.elabscience.com)

Tel: 1-832-243-6086  
Email: [techsupport@elabscience.com](mailto:techsupport@elabscience.com)

Fax: 1-832-243-6017

Rev. V1.0

Immunofluorescent analysis of (100% Ice-cold methanol)  
fixed LNCaP cells using anti-Na<sup>+</sup>/K<sup>+</sup> ATPase alpha-1  
Monoclonal Antibody at dilution of 1:50.

## Preparation & Storage

**Storage** Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.  
**Shipping** Ice bag

## Background

The protein encoded by this gene belongs to the family of P-type cation transport ATPases, and to the subfamily of Na<sup>+</sup>/K<sup>+</sup> -ATPases. Na<sup>+</sup>/K<sup>+</sup> -ATPase is an integral membrane protein responsible for establishing and maintaining the electrochemical gradients of Na and K ions across the plasma membrane. These gradients are essential for osmoregulation, for sodium-coupled transport of a variety of organic and inorganic molecules, and for electrical excitability of nerve and muscle. The catalytic subunit of Na<sup>+</sup>/K<sup>+</sup> -ATPase is encoded by multiple genes. This is the catalytic component of the active enzyme, which catalyzes the hydrolysis of ATP coupled with the exchange of sodium and potassium ions across the plasma membrane. This action creates the electrochemical gradient of sodium and potassium ions, providing the energy for active transport of various nutrients.

## For Research Use Only

Toll-free: 1-888-852-8623  
Web: [www.elabscience.com](http://www.elabscience.com)

Tel: 1-832-243-6086  
Email: [techsupport@elabscience.com](mailto:techsupport@elabscience.com)

Fax: 1-832-243-6017

Rev. V1.0