Elabscience®


Elab Bright[™] Violet 421 Anti-Mouse CD4 Antibody[GK1.5]

Catalog Number: E-AB-F1097Q2

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description	
Reactivity	Mouse
Host	Rat
lsotype	Rat lgG2b, κ
Clone No.	GK1.5
Isotype Control	Elab Bright™ Violet 421 Rat IgG2b, κ Isotype Control[R35-38] [Product AN00821Q2]
Conjugation	Elab Bright™Violet 421
Conjugation Information	Elab Bright Violet 421 is designed to be excited by the violet laser (405 nm) and detected using an optical filter centered near 421 nm (e.g., a 450/50 nm bandpass filter).
Storage Buffer	Phosphate buffered solution, pH 7.2, containing 0.09% sodium azide and 1% BSA.
Applications	Recommended usage
FCM	Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5 μ L of antibody per test (million cells in 100 μ L staining volume or per 100 μ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.

Data



Staining of C57BL/6 murine splenocytes cells with APC Anti-Mouse CD3 Antibody and Elab Bright Violet 421 Rat Anti-Mouse CD4 Antibody[GK1.5] (left) or Elab Bright Violet 421 Rat IgG2b, κ Isotype Control (right). Total viable cells were used for analysis.

Preparation & Storag	ye
Storage	Keep as concentrated solution.
	This product can be stored at 2-8°C for 12 months. Please protected from prolonged exposure to light and do not freeze.
Shipping	Ice bag
Antigen Information	
Alternate Names	L3T4;T-cell surface antigen T4/Leu-3;T-cell surface glycoprotein CD4;T4
Uniprot ID	P06332
Gene ID	12504

Elabscience®

Elabscience Biotechnology Co., Ltd. A Reliable Research Partner in Life Science and Medicine

Background

CD4 is a 55 kD protein also known as L3T4 or T4. It is a member of the Ig superfamily, primarily expressed on most thymocytes, a subset of T cells, and weakly on macrophages and dendritic cells. It acts as a coreceptor with the TCR during T cell activation and thymic differentiation by binding MHC class II and associating with the protein tyrosin kinase, Ick.