

APC/Cyanine 7 Anti-Human CD154 (CD40L) Antibody[24-31]

Catalog Number: AN00341N

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description


Reactivity	Human
Host	Mouse
Isotype	Mouse IgG1, κ
Clone No.	24-31
Isotype Control	APC/Cyanine 7 Mouse IgG1, κ Isotype Control[MOPC-21] [Product E-AB-F09792N]
Conjugation	APC/Cyanine 7
Conjugation Information	APC/Cyanine 7 is designed to be excited by the Red (627-640 nm) lasers and detected using an optical filter centered near 780 nm (e.g., a 780/60 nm bandpass filter).
Storage Buffer	Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer.

Applications

Recommended usage

FCM	Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5 μL of antibody per test (million cells in 100 μL staining volume or per 100 μL of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.
------------	---

Data



Staining of MIX stimulated(5h) PBMC with APC/Cyanine 7 Anti-Human CD154 Antibody[24-31](filled gray histogram) or APC/Cyanine 7 Mouse IgG1, κ Isotype Control(empty black histogram).

Preparation & Storage

Storage	Keep as concentrated solution. This product can be stored at 2-8°C for 24 months. Please protected from prolonged exposure to light and do not freeze.
Shipping	Ice bag

Antigen Information

Alternate Names	CD40L;gp39;TRAP;T-BAM;TNFSF5
Uniprot ID	P29965
Gene ID	959

For Research Use Only

Toll-free: 1-888-852-8623
Web: www.elabscience.com

Tel: 1-832-243-6086
Email: techsupport@elabscience.com

Fax: 1-832-243-6017

Rev. V1.2

Background

CD154 (CD40 ligand) is also known as CD40L, gp39, TRAP and T-BAM. CD40 ligand is a 32-39 kD type II transmembrane glycoprotein. It is a member of the TNF superfamily and is expressed on activated T cells. It has been reported to be important for B cell costimulation following binding of its receptor, CD40. Additionally, binding of CD40L to CD40 on B cells promotes the secretion of immunoglobulin and Ig isotype switching. CD40L is also involved in the regulation of cytokine production by T cells.