## **Recombinant VAPB/VAP-B Monoclonal Antibody**

## catalog number: AN300203P

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Ice bag

Description	
Reactivity	Human
Immunogen	Recombinant Human VAPB / VAP-B protein
Host	Rabbit
Isotype	IgG
Clone	8C6
Purification	Protein A
Buffer	0.2 µm filtered solution in PBS
Applications	Recommended Dilution
WB	1:500-1:2000
Data	
dilution of 1:500. Lane B: A549 Whole Cell I Lysate, Lysate <b>Obse</b>	B / VAP-B Monoclonal Antibody at A: HepG2 Whole Cell Lysate, Lane Lysate, Lane C: Jurkat Whole Cell s/proteins at 30 μg per lane. rved-MW:32 kDa
dilution of 1:500. Lane B: A549 Whole Cell I Lysate, Lysate <b>Obse</b> <b>Calcu</b>	A: HepG2 Whole Cell Lysate, Lane Lysate, Lane C: Jurkat Whole Cell s/proteins at 30 μg per lane. rved-MW:32 kDa lated-MW:27 kDa
dilution of 1:500. Lane B: A549 Whole Cell I Lysate, Lysate Obse Calcu Preparation & Storage	A: HepG2 Whole Cell Lysate, Lane Lysate, Lane C: Jurkat Whole Cell s/proteins at 30 μg per lane. rved-MW:32 kDa lated-MW:27 kDa
dilution of 1:500. Lane B: A549 Whole Cell I Lysate, Lysate <b>Obse</b> <b>Calcu</b>	A: HepG2 Whole Cell Lysate, Lane Lysate, Lane C: Jurkat Whole Cell s/proteins at 30 μg per lane. rved-MW:32 kDa lated-MW:27 kDa

Shipping Background

The protein encoded by this gene is a type IV membrane protein found in plasma and intracellular vesicle membranes. The encoded protein is found as a homodimer and as a heterodimer with VAPA. This protein also can interact with VAMP1 and VAMP1 and may be involved in vesicle trafficking.