

Apolipoprotein H/APOH Monoclonal Antibody

catalog number: **AN200209P**

Note: *Centrifuge before opening to ensure complete recovery of vial contents.*


Description

Reactivity	Human
Immunogen	Recombinant Human APOH protein
Host	Mouse
Isotype	IgG1
Clone	12F6
Purification	Protein A
Buffer	0.2 µm filtered solution in PBS


Applications Recommended Dilution

ICC/IF	1:20-1:100
FCM	1:25-1:100

Data



Immunofluorescence analysis of Human APOH in HepG2 or Hela cells. Cells were fixed with 4% PFA, permeabilized with 1% Triton X-100 in PBS, blocked with 10% serum, and incubated with mouse anti-Human APOH monoclonal antibody (1:60). Then cells were stained with the Alexa Fluor® 488-conjugated Goat Anti-mouse IgG secondary antibody (left panel, captured by laser confocal scanning microscope; right panel, captured by fluorescence microscope), counterstained with DAPI (blue). Positive staining was localized to cytoplasm.



Flow cytometric analysis of Human APOH expression on MDA-MB-231 cells. The cells were treated according to manufacturer's manual, stained with purified anti-Human APOH, then a FITC-conjugated second step antibody. The fluorescence histograms were derived from gated events with the forward and side light-scatter characteristics of intact cells.

Preparation & Storage

Storage	This antibody can be stored at 2°C-8°C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20°C to -80°C. Preservative-Free. Avoid repeated freeze-thaw cycles.
Shipping	Ice bag

Background

Apolipoprotein H has been implicated in a variety of physiologic pathways including lipoprotein metabolism, coagulation, and the production of antiphospholipid autoantibodies. APOH may be a required cofactor for anionic phospholipid binding by the antiphospholipid autoantibodies found in sera of many patients with lupus and primary antiphospholipid syndrome, but it does not seem to be required for the reactivity of antiphospholipid autoantibodies associated with infections.

For Research Use Only