

## Human HAVCR2 Antibody Pair Set

<b>Catalog No.</b>	E-KAB-0437	<b>Applications</b>	ELISA
<b>Synonyms</b>	T cell immunoglobulin and mucin domain containing protein 3; hepatitis virus cellular receptor 2; CD366		

### Kit components & Storage

Title	Specifications	Storage
Human HAVCR2 Capture Antibody	1 vial, 100 µg	Store at -20℃ for one year. Avoid freeze/thaw cycles.
Human HAVCR2 Detection Antibody (Biotin)	1 vial, 50 µL	Store at -20℃ for one year. Avoid freeze/thaw cycles.

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


### Product Information

Items		Characteristic (E-KAB-0437)	
		Human HAVCR2 Capture Antibody	Human HAVCR2 Detection Antibody (Biotin)
Immunogen Information	Immunogen	Recombinant Human HAVCR2 protien	Recombinant Human HAVCR2 protien
	Swissprot	Q8TDQ0	
Product details	Reactivity	Human	Human
	Host	Rabbit	Rabbit
	Conjugation	Unconjugated	Biotin
	Concentration	0.5 mg/mL	/
	Buffer	PBS with 0.04% Proclin 300; 50% glycerol; pH 7.5	PBS with 0.04% Proclin 300; 1% protective protein; 50% glycerol; pH 7.5
	Purify	Protein A & Antigen Affinity	Protein A
	Specificity	Detects Human HAVCR2 in ELISAs.	

### For Research Use Only

## Applications

### Human HAVCR2 Sandwich ELISA Assay

	Recommended Concentration/Dilution	Reagent	Images
ELISA Capture	0.5-4 µg/mL	Human HAVCR2 Capture Antibody	
ELISA Detection	1:1000-1:10000	Human HAVCR2 Detection Antibody (Biotin)	

**Note:** This standard curve is only for demonstration purposes. A standard curve should be generated for each assay!

## Background

T cell immunoglobulin and mucin domain-3 (TIM3), also called hepatitis A virus cellular receptor 2 (HAVCR2), is a transmembrane glycoprotein of the TIM family of immune regulating molecules and plays an important role in the Th1-mediated immune response. TIM3 is expressed on the Th1 cells, CD8 T-cells, monocytes, and dendritic cells, but not on Th2 cells. TIM3 expressed by monocytes and dendritic cells facilitates phagocytosis of apoptotic cells and up-regulates cross-presentation of apoptotic cell-associated antigens through interaction with phosphatidylserine. Engagement of TIM3 by its ligand galectin-9 induces a range of immunosuppressive functions which enhance immune tolerance and inhibit anti-tumor immunity. Stimulation of TIM3 with an agonistic antibody promotes inflammation through the activation of innate immune cells. TIM3 is also regarded as a potential target molecule for immunotherapy. TIM3 and programmed cell death 1 (PD-1) as two important coinhibitory regulators of T cell responses, have been implicated with the T-cell dysfunction or exhaustion associated with chronic HBV infection including HBV-related HCC.

## For Research Use Only